

IWASAWA THEORY OF TOTALLY REAL FIELDS FOR CERTAIN NON-COMMUTATIVE p -EXTENSIONS

TAKASHI HARA

ABSTRACT. In this paper, we prove the Iwasawa main conjecture (in the sense of [CFKSV]) of totally real fields for certain specific non-commutative p -adic Lie extensions, using the integral logarithms introduced by Oliver and Taylor. Our result gives certain generalization of Kato's proof of the main conjecture for Galois extensions of Heisenberg type ([Kato1]).

0. INTRODUCTION

0.1. Introduction. The Iwasawa main conjecture, which describes mysterious relation between “arithmetic” characteristic elements and “analytic” p -adic zeta functions, has been proven under many situations for abelian extensions. However, for non-abelian extensions, it took many years even to formulate the main conjecture. In 2005, Coates et al. formulated it ([CFKSV]) by using algebraic K -theory (especially the localization exact sequence), and Kazuya Kato has proven it for certain specific p -adic Lie extensions of totally real fields up to the present ([Kato1]). Mahesh Kakde generalized Kato's proof and proved the main conjecture for other types of extensions ([Kakde]). Jürgen Ritter and Alfred Weiss also formulated the main conjecture in a little different way (“Equivariant Iwasawa theory,” see [R-W1].), also using algebraic K -theory.

In this paper, we prove the Iwasawa main conjecture (in the sense of [CFKSV]) of totally real fields for certain non-commutative p -extensions, using the method of Kato in [Kato1].

Let F be a totally real number field, and let p be a prime number. Let F^∞/F be a totally real Lie extension containing the cyclotomic $\mathbb{Z}_p$ -extension F^{cyc} of F . We assume that only finitely many primes of F ramify in F^∞ .

The aim of this paper is to prove the following theorem.

Theorem 0.1 (=Theorem 3.1). *Assume that the following conditions are satisfied.*

$$(1) \quad G = \text{Gal}(F^\infty/F) \cong \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \Gamma,$$

where Γ is a commutative p -adic Lie group isomorphic to $\mathbb{Z}_p$.

(2) $p \neq 2, 3$.

- (3) *There exists a finite extension $F' \subseteq F^\infty$ of F such that the μ -invariant of $(F')^{\text{cyc}}/F'$ equals to zero, where $(F')^{\text{cyc}}/F'$ is the cyclotomic $\mathbb{Z}_p$ -extension of F' .*

Then the p -adic zeta function $\xi_{F^\infty/F}$ for F^∞/F exists and the Iwasawa main conjecture for F^∞/F is true.

Let us summarize how to prove this theorem. We consider the following family of subgroups of G .

$$\begin{aligned}
U_0 &= G, & V_0 &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \{1\}, \\
U_1 &= \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \Gamma, & V_1 &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \{1\}, \\
\widetilde{U}_2 &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \Gamma, & \widetilde{V}_2 &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \{1\}, \\
U_2 &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \Gamma, & V_2 &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \{1\}, \\
U_3 &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \Gamma, & V_3 &= \{I_4\} \times \{1\},
\end{aligned}$$

where I_4 is the unit matrix of $\text{GL}_4(\mathbb{F}_p)$.

In the following, we use the notation U_i (resp. V_i) for one of the subgroups $U_0, U_1, \widetilde{U}_2, U_2$ and U_3 (resp. $V_0, V_1, \widetilde{V}_2, V_2$ and V_3). Note that each quotient group U_i/V_i is abelian.

Now we have a homomorphism

$$\theta_i: K_1(\Lambda(G)) \xrightarrow{\text{Nr}} K_1(\Lambda(U_i)) \rightarrow K_1(\Lambda(U_i/V_i)) = \Lambda(U_i/V_i)^\times$$

where $\Lambda(G)$ denotes the Iwasawa algebra of G . Here the first map is the norm map of K -theory and the second one is the natural map induced by $\Lambda(U_i) \rightarrow \Lambda(U_i/V_i)$.

On the other hand, John Coates et al. introduced the canonical Ore set S of $\Lambda(G)$ (See §2 and [CFKSV]) and considered the Ore localization $\Lambda(G)_S$ to formulate the main conjecture. For this localized Iwasawa algebra, we also have a homomorphism

$$\theta_{S,i}: K_1(\Lambda(G)_S) \xrightarrow{\text{Nr}} K_1(\Lambda(U_i)_S) \rightarrow K_1(\Lambda(U_i/V_i)_S) = \Lambda(U_i/V_i)_S^\times$$

by the same construction as θ_i . We denote

$$\theta = (\theta_i)_i: K_1(\Lambda(G)) \rightarrow \prod_i \Lambda(U_i/V_i)^\times$$

and

$$\theta_S = (\theta_{S,i})_i: K_1(\Lambda(G)_S) \rightarrow \prod_i \Lambda(U_i/V_i)_S^\times.$$

Then we have the following proposition.

Proposition 0.2 (Proposition 5.4, Proposition 6.3 and Proposition 6.4).
There exist subgroups

$$\Psi \subseteq \prod_i \Lambda(U_i/V_i)^\times$$

and

$$\Psi_S \subseteq \prod_i \Lambda(U_i/V_i)_S^\times$$

which satisfy the following conditions:

- (1) $\text{Image}(\theta_S) \subseteq \Psi_S$.
- (2) $\text{Image}(\theta) = \Psi$.
- (3) $\Psi_S \cap \prod_i \Lambda(U_i/V_i)^\times = \Psi$.

We can characterize both Ψ and Ψ_S as the subgroups consisting of all elements which satisfy certain norm relations and certain congruences (for details, see Definition 5.3 and Proposition 6.2.). In the following, we denote the induced homomorphisms by the same symbols

$$\theta: K_1(\Lambda(G)) \rightarrow \Psi \quad \text{and} \quad \theta_S: K_1(\Lambda(G)_S) \rightarrow \Psi_S.$$

We call the surjective homomorphism θ the *theta map* for G and the homomorphism θ_S the *localized theta map* for G .

Then we obtain the following outstanding theorem which was first observed by David Burns.

Theorem 0.3 (=Theorem 3.4, Burns). *Let ξ_i be the p -adic zeta function for the abelian p -adic Lie extension F_{V_i}/F_{U_i} where F_{U_i} (resp. F_{V_i}) is the maximal intermediate field of F^∞/F fixed by U_i (resp. V_i). If $(\xi_i)_i$ is contained in Ψ_S , the p -adic zeta function ξ for F^∞/F exists as an element of $K_1(\Lambda(G)_S)$ and satisfies the main conjecture.*

Note that the p -adic zeta function (pseudomeasure) ξ_i for F_{V_i}/F_{U_i} has been constructed by using the theory of Pierre R. Deligne and Kenneth A. Ribet ([De-Ri]), and the Iwasawa main conjecture for F_{V_i}/F_{U_i} has been proven by Andrew Wiles ([Wiles]).

The condition for $(\xi_i)_i$ to be contained in Ψ_S is essentially given by the congruences among ξ_i 's, so we may reduce the non-commutative Iwasawa main conjecture to the congruences among the abelian p -adic zeta pseudomeasures via the theta map.

In order to study the congruences which abelian p -adic zeta pseudomeasures should satisfy, we use the theory of Hilbert modular forms of Deligne-Ribet ([De-Ri]), and derive the congruences of p -adic zeta pseudomeasures (that is, the congruences of constant terms of certain Λ -adic Hilbert modular

forms) from those of the coefficients of non-constant terms of Λ -adic Hilbert modular forms (See §7). Kato and Ritter-Weiss first used this technique in [Kato1] and [R-W3], and obtained many kinds of congruences among abelian p -adic zeta pseudomeasures.

Actually it is difficult to prove all the desired congruences by using only this technique, therefore we use the existence of the p -adic zeta function for a certain quotient group $\overline{G}$ of G , proven by Kato in [Kato1]. We prove our main theorem (Theorem 0.1) by using an inductive technique.

0.2. Overview. The detailed content of this paper is as follows.

In §1, we review basic results of (classical) algebraic K -theory. In particular, we summarize properties of integral logarithmic homomorphisms, which were first introduced by Robert Oliver and Laurence R. Taylor to study the structure of the K_1 -group of a group ring $R[G]$ where G is a finite group and R is the integer ring of a finite extension of $\mathbb{Q}_p$. The integral logarithmic homomorphisms are maps from multiplicative K_1 -groups to certain additive groups, which are much easier to treat (Proposition-Definition 1.29).

In §2, we review the theory of Coates et al. ([CFKSV]), especially how to formulate the main conjecture.

We state our main theorem precisely in §3, and introduce Burns' technique under more general situations than Theorem 0.1.

We construct the theta map for our case from §4 to §6. In §4, we construct the additive version of the theta map (Proposition-Definition 4.3) by using linear representation theory of finite groups. In §5, we translate the results on the additive theta map proven in §4 into the (multiplicative) theta map, using the integral logarithmic homomorphisms. In §6, we construct the localized version of the theta map θ_S . For this purpose, we take the p -adic completion $\widehat{\Lambda(\Gamma)_{(p)}}$ of $\Lambda(\Gamma)_{(p)}$, and apply the arguments of §4 and §5 to $\widehat{\Lambda(\Gamma)_{(p)}}[G^f]$.

The condition for abelian p -adic zeta pseudomeasures to be contained in Ψ_S is essentially given as the congruences among them. Hence in §7, we study congruences which abelian p -adic zeta pseudomeasures satisfy (Proposition 7.2). We use the theory of Deligne and Ribet on Hilbert modular forms ([De-Ri]).

Since the congruences obtained in §7 are not sufficient to conclude that abelian p -adic zeta pseudomeasures are contained in Ψ_S , we introduce Kato's p -adic zeta function for a certain sub p -adic Lie extension F_N/F of F^∞/F (Theorem 8.1), and prove Theorem 0.1 by an inductive technique.

Our main theorem gives a new example which is not deduced from previous results ([Kato1], [R-W2], [R-W3], and [R-W4]).

CONTENTS

0. Introduction	1
1. Preliminaries on algebraic K -theory	6
2. Basic results of non-commutative Iwasawa theory	18
3. The main theorem and Burns' technique	23
4. The additive theta map	28
5. Translation into the multiplicative theta map	39

6. Localized theta map	56
7. Congruences among abelian p -adic zeta pseudomeasures	60
8. Proof of the main theorem	72
References	76

0.3. Notation. In this paper, p always denotes a positive prime number.

We denote by $\mathbb{N}$ the set of natural numbers (the set of *strictly* positive integers), denote by $\mathbb{Z}$ (resp. $\mathbb{Z}_p$) the ring of integers (resp. p -adic integers), and also denote by $\mathbb{Q}$ (resp. $\mathbb{Q}_p$) the rational number field (resp. the p -adic number field).

For an arbitrary group G , we denote by $\text{Conj}(G)$ the set of all conjugacy classes of G .

For every pro-finite group P , we always denote by $\Lambda(P) = \mathbb{Z}_p[[P]]$ its Iwasawa algebra (that is, its completed group ring over $\mathbb{Z}_p$).

We denote by Γ a commutative p -adic Lie group which is isomorphic to $\mathbb{Z}_p$. Throughout this paper, we fix a topological generator t of Γ . In other words, we fix an isomorphism

$$\begin{aligned} \Lambda(\Gamma) &\xrightarrow{\sim} \mathbb{Z}_p[[T]] \\ t &\mapsto 1 + T \end{aligned}$$

where $\mathbb{Z}_p[[T]]$ is the formal power series ring over $\mathbb{Z}_p$.

We always assume that every associative ring has 1. We also use the following notation:

$$\begin{aligned} M_n(R) &= \{\text{the ring of } n \times n\text{-matrices with entries in } R\}, \\ GL_n(R) &= \{M \in M_n(R) \mid M \text{ is an invertible matrix}\}, \end{aligned}$$

where R is an associative ring. If R is a commutative domain, we denote by $\text{Frac}(R)$ its fractional field.

We always regard K_0 -groups as additive groups, and K_1 -groups as multiplicative groups.

Finally, we fix an algebraic closure $\overline{\mathbb{Q}}$ of $\mathbb{Q}$ and fix embeddings

$$\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}, \quad \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}_p}$$

where $\mathbb{C}$ denotes the complex number field and $\overline{\mathbb{Q}_p}$ the algebraic closure of $\mathbb{Q}_p$.

0.4. Acknowledgment. The author would like to thank everyone with whom he has been concerned. He would like to express his sincere gratitude to Professor Shuji Saito, Professor Kazuya Kato and Professor Takeshi Tsuji among them; Professor Shuji Saito has invited the author to the mysterious and interesting world of number theory through his lectures and seminars; Professor Kazuya Kato has shown the author the mystic aspect of non-commutative Iwasawa theory and motivated the author to study it through his intensive lectures at the University of Tokyo in 2006; and Professor Takeshi Tsuji, whom the author is most grateful to, has given the author a lot of useful advice through his seminars and read the manuscript very carefully. Especially, the main idea of the inductive technique (used in

§8) is due to him. This paper could never have existed without their direct and indirect cooperation.

Mahesh Kakde has recently generalized the Kato's method used in [Kato1] and proven the main conjecture for other cases in the different way from this paper (see [Kakde]). The author would like to express his sincere gratitude to Mahesh Kakde for sending the latest version of his paper.

1. PRELIMINARIES ON ALGEBRAIC K -THEORY

In this section, we summarize basic results of algebraic K -theory which we need to formulate the non-commutative Iwasawa main conjecture and to construct the theta map.

1.1. Definitions and first properties. First, we review the definition of K -groups and their properties. For more details, see [Bass1].

Definition 1.1 (Grothendieck groups, K_0 -groups). Let $\mathcal{C}$ be a category with a product $\perp$ (recall that a category $\mathcal{C}$ with a product $\perp$ is a category equipped with a functor $\perp: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$). Then we define *the Grothendieck group of $\mathcal{C}$* $K_0(\mathcal{C})$ as an abelian group equipped with a map

$$[\cdot]: \text{Ob } \mathcal{C} \longrightarrow K_0(\mathcal{C})$$

satisfying the following universal properties.

(K_0 -1) For every $X, Y \in \text{Ob } \mathcal{C}$ satisfying $X \cong Y$, $[X] = [Y]$.

(K_0 -2) For every $X, Y \in \text{Ob } \mathcal{C}$, $[X \perp Y] = [X] + [Y]$.

Namely, if a map $f: \text{Ob } \mathcal{C} \longrightarrow A$ (A : an abelian group) satisfies the properties (K_0 -1) and (K_0 -2), there exists a unique group homomorphism $\psi: K_0(\mathcal{C}) \rightarrow A$ which makes the following diagram commute:

$$\begin{array}{ccc} \text{Ob } \mathcal{C} & \xrightarrow{[\cdot]} & K_0(\mathcal{C}) \\ & \searrow f & \downarrow \exists! \psi \\ & & A \end{array}$$

For every associative ring R , we define $K_0(R)$ as the Grothendieck group of the category of finitely generated projective left R -modules.

Definition 1.2 (Whitehead groups, K_1 -groups). Let $\mathcal{C}$ be a category with a product $\perp$. Let $\text{Aut}(\mathcal{C})$ be the category of automorphisms of objects of $\mathcal{C}$. Namely, an object of $\text{Aut}(\mathcal{C})$ is a pair (X, σ) where $X \in \text{Ob } \mathcal{C}$ and $\sigma: X \rightarrow X$ is an automorphism of X . A morphism $f: (X, \sigma) \rightarrow (Y, \tau)$ is a morphism $f: X \rightarrow Y$ in $\mathcal{C}$ which satisfies $f \circ \sigma = \tau \circ f$.

Then we define *the Whitehead group of $\mathcal{C}$* $K_1(\mathcal{C})$ as an abelian group equipped with a map

$$[\cdot]: \text{Ob } \text{Aut}(\mathcal{C}) \longrightarrow K_1(\mathcal{C})$$

satisfying the following universal properties.

(K_1 -1) For every $(X, \sigma), (Y, \tau) \in \text{Ob } \text{Aut}(\mathcal{C})$ satisfying $(X, \sigma) \cong (Y, \tau)$, $[(X, \sigma)] = [(Y, \tau)]$.

(K_1 -2) For every $(X, \sigma), (Y, \tau) \in \text{Ob } \text{Aut}(\mathcal{C})$, $[(X, \sigma) \perp (Y, \tau)] = [(X, \sigma)] \cdot [(Y, \tau)]$.

(K_1 -3) For every $(X, \sigma), (X, \sigma') \in \text{Ob Aut}(\mathcal{C})$, $[(X, \sigma \circ \sigma')] = [(X, \sigma)] \cdot [(X, \sigma')]$.

For every associative ring R , we define $K_1(R)$ as the Whitehead group of the category of finitely generated projective left R -modules.

Remark 1.3. When R is the group ring $\mathcal{O}_K[G]$ of a finite group G over the integer ring of a number field K , the terminology “Whitehead group” is usually used for the group

$$\text{Wh}(\mathcal{O}_K[G]) = K_1(\mathcal{O}_K[G]) / (\mu(\mathcal{O}_K) \times G^{\text{ab}}),$$

where $\mu(\mathcal{O}_K)$ is the multiplicative group consisting of all roots of 1 contained in $\mathcal{O}_K$.

We have another interpretation of the Whitehead group of an associative ring R (Whitehead’s construction): let $E_n(R)$ be a subgroup of $GL_n(R)$ generated by all “elementary matrices,” that is,

$$E_n(R) = \langle I_n + rE_{ij} \mid 1 \leq i \neq j \leq n, r \in R \rangle$$

where

$$E_{ij} = \begin{matrix} & & & j & \\ i & \begin{pmatrix} 0 & 0 & \cdots & & 0 \\ 0 & \ddots & & & 1 \\ & & \ddots & & \\ \vdots & & & \ddots & \vdots \\ 0 & \cdots & & \cdots & 0 \end{pmatrix} \end{matrix}.$$

Here we denote the unit matrix of $GL_n(R)$ by I_n . Note that $E_n(R)$ is normal in $GL_n(R)$.

Let

$$GL(R) = \varinjlim_n GL_n(R) \quad \text{and} \quad E(R) = \varinjlim_n E_n(R),$$

then we have

$$K_1(R) = GL(R)/E(R).$$

For the equivalence of Definition 1.2 and Whitehead construction, see [Bass1], Chapter IX.

Definition 1.4 (Relative Whitehead groups). Let R be an associative ring and $\mathfrak{a} \subseteq R$ an arbitrary (two-sided) ideal. Set

$$\begin{aligned} GL_n(R, \mathfrak{a}) &= \text{Ker}(GL_n(R) \longrightarrow GL_n(R/\mathfrak{a})), \\ E_n(R, \mathfrak{a}) &= \text{The minimal normal subgroup of } GL_n(R, \mathfrak{a}) \\ &\quad \text{containing } \{I_n + rE_{ij} \mid 0 \leq i \neq j \leq n, r \in \mathfrak{a}\}, \end{aligned}$$

and

$$GL(R, \mathfrak{a}) = \varinjlim_n GL_n(R, \mathfrak{a}), \quad E(R, \mathfrak{a}) = \varinjlim_n E_n(R, \mathfrak{a}).$$

Then we define

$$K_1(R, \mathfrak{a}) = GL(R, \mathfrak{a})/E(R, \mathfrak{a}).$$

Proposition 1.5 (Whitehead's lemma). *Let R and $\mathfrak{a}$ be as above. Then*

$$E(R) = [\mathrm{GL}(R), \mathrm{GL}(R)],$$

$$E(R, \mathfrak{a}) = [E(R), E(R, \mathfrak{a})] = [\mathrm{GL}(R), \mathrm{GL}(R, \mathfrak{a})].$$

Proof. See [Milnor], Lemma 3.1, Lemma 4.3. $\square$

The following two propositions are well known and we often use them later.

Proposition 1.6. *Let R be an associative ring and $\mathfrak{a}$ be a two-sided ideal contained in its Jacobson radical. Then the canonical homomorphism*

$$K_1(R) \longrightarrow K_1(R/\mathfrak{a})$$

induced by $R \rightarrow R/\mathfrak{a}$ is surjective.

Proof. See [Bass1], Chapter IX, Proposition (1.3). $\square$

Proposition 1.7. *Let R be a semi-local associative ring (recall that R is semi-local if R/J is semi-simple where J is the Jacobson radical of R). Then the following properties hold.*

(1) *The group homomorphism*

$$R^\times \longrightarrow K_1(R); u \mapsto [(R, - \cdot u)]$$

is surjective, where $- \cdot u$ is the automorphism of R defined by multiplication by u from the right (here we regard R as a left R -module).

If R is semi-local and also commutative, the homomorphism above is an isomorphism.

(2) *(stability property)*

For each $d \geq 2$, we have the canonical isomorphism

$$K_1(R) \cong \mathrm{GL}_d(R)/E_d(R).$$

Proof. For (1), see [Bass1] Chapter IX, Proposition (1.4). When R is commutative, the determinant map gives the inverse map of $R^\times \rightarrow K_1(R)$.

For (2), see [Bass1] Chapter V, Theorem (9.1). $\square$

Now let us study the project limit of K_1 -groups for semi-local rings.

Proposition 1.8. *Let R be a semi-local ring and $\{R^{(n)}\}_{n \in \mathbb{N}}$ a projective system of semi-local rings such that $R^{(n)} \rightarrow R^{(m)}$ is surjective for each $n > m \geq 1$ and $\varprojlim_n R^{(n)} \cong R$.*

Then the homomorphism

$$K_1(R) \rightarrow \varprojlim_n K_1(R^{(n)})$$

induced by the canonical homomorphisms $K_1(R) \rightarrow K_1(R^{(n)})$ is an isomorphism.

Proof. By the stability property (Proposition 1.7 (2)), we have

$$K_1(R) \cong \mathrm{GL}_d(R)/E_d(R) \quad \text{and} \quad K_1(R^{(n)}) \cong \mathrm{GL}_d(R^{(n)})/E_d(R^{(n)})$$

for every $d \geq 2$. Fix $d \geq 2$.

Consider the exact sequence

$$(1.1) \quad 1 \longrightarrow E_d(R^{(n)}) \longrightarrow \mathrm{GL}_d(R^{(n)}) \longrightarrow K_1(R^{(n)}) \longrightarrow 1$$

for each $n \geq 1$. Then for every $n > m \geq 1$, the natural homomorphism

$$E_d(R^{(n)}) \rightarrow E_d(R^{(m)})$$

induced by $R^{(n)} \rightarrow R^{(m)}$ is clearly surjective. Therefore $\{E_d(R^{(n)})\}_{n \in \mathbb{N}}$ satisfies the Mittag-Leffler condition, and we have the following exact sequence

$$1 \longrightarrow \varprojlim_n E_d(R^{(n)}) \longrightarrow \varprojlim_n GL_d(R^{(n)}) \longrightarrow \varprojlim_n K_1(R^{(n)}) \longrightarrow 1$$

by taking the projective limit of (1.1).

Now consider the following diagram:

$$\begin{array}{ccccccccc} 1 & \longrightarrow & E_d(R) & \longrightarrow & GL_d(R) & \longrightarrow & K_1(R) & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \varprojlim_n E_d(R^{(n)}) & \longrightarrow & \varprojlim_n GL_d(R^{(n)}) & \longrightarrow & \varprojlim_n K_1(R^{(n)}) & \longrightarrow & 1 \end{array}$$

here the vertical homomorphisms are induced by the canonical homomorphism $R \rightarrow R^{(n)}$. We can easily check that each square diagram commutes.

Then the left and middle vertical homomorphisms are canonically isomorphic. Hence the right vertical homomorphism is also an isomorphism by the snake lemma. $\square$

1.2. Norm maps in K -theory. The theta map, which we will construct in the following sections, is essentially a family of norm maps in algebraic K -theory.¹ So let us review the construction and properties of norm maps of K -groups.

Let R'/R be an extension of a ring R . Suppose that R' is a finitely generated projective module as a left R -module. Then we may regard R' as left R -right R' -bimodule ${}_R R'_{R'}$. We define

$$\mathrm{Nr}_{R'/R} := [{}_R R'_{R'} \otimes_{R'} -] : K_i(R') \longrightarrow K_i(R) \quad (i = 0, 1).$$

We often use norm maps in the following situation: let G be a group and H be its subgroup of finite index. Then we have an inclusion of group rings $\mathbb{Z}_p[H] \rightarrow \mathbb{Z}_p[G]$. Hence we obtain a norm map

$$\mathrm{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} : K_i(\mathbb{Z}_p[G]) \longrightarrow K_i(\mathbb{Z}_p[H]) \quad (i = 0, 1).$$

If G is a pro-finite group and H is its open subgroup, we also obtain $\mathrm{Nr}_{\Lambda(G)/\Lambda(H)}$ by the same construction.

Now we calculate

$$\mathrm{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} : K_1(\mathbb{Z}_p[G]) \longrightarrow K_1(\mathbb{Z}_p[H]) \cong \mathbb{Z}_p[H]^\times$$

under the specific condition that G is a group and H is a commutative subgroup of finite index. Note that both $\mathbb{Z}_p[G]$ and $\mathbb{Z}_p[H]$ are local rings. By Proposition 1.7 (1), we can identify $K_1(\mathbb{Z}_p[H])$ with $\mathbb{Z}_p[H]^\times$. Take a system of representatives $\{u_1, \dots, u_r\}$ of the coset decomposition $H \backslash G$. Then $\mathbb{Z}_p[G]$ is regarded as a left free $\mathbb{Z}_p[H]$ -module with basis $\{u_1, \dots, u_r\}$.

¹In some books and papers, norm maps are also called “transfer homomorphisms.”

Let ϕ be an element of $K_1(\mathbb{Z}_p[G])$. By Proposition 1.7 (1) again, we obtain $x \in \mathbb{Z}_p[G]^\times$ such that $[x] = \phi$. Let

$$u_j x = \sum_{i=1}^r x_{ij} u_i \quad (x_{ij} \in \mathbb{Z}_p[H])$$

for each j . Then we can calculate $\text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} \phi$ as

$$\text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} \phi = \det((x_{ij})_{1 \leq i, j \leq r}) \in \mathbb{Z}_p[H]^\times.$$

Proof. By the calculation above, we may identify $\text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} \phi$ with the image of $(x_{ij})_{i,j}$ in $K_1(\mathbb{Z}_p[H]) = \text{GL}(\mathbb{Z}_p[H])/\text{E}(\mathbb{Z}_p[H])$. Since $\mathbb{Z}_p[H]$ is commutative, the surjection $\mathbb{Z}_p[H]^\times \rightarrow K_1(\mathbb{Z}_p[H])$ is an isomorphism by Proposition 1.7 (1). The determinant map gives the inverse map of the isomorphism above, and it maps $\text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} \phi = [(x_{ij})_{i,j}]$ to $\det(x_{ij})_{i,j}$. $\square$

When G is a pro-finite group and H is its commutative open subgroup, we may calculate $\text{Nr}_{\Lambda(G)/\Lambda(H)}$ explicitly in the same way.

We end this subsection with a certain compatibility property of norm maps.

Proposition 1.9. *Let G be a group and H be its subgroup of finite index. Let N be a subgroup of H normal in G and H .*

Then for $i = 0, 1$, the following diagram commutes:

$$\begin{array}{ccc} K_i(\mathbb{Z}_p[G]) & \xrightarrow{\text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]}} & K_i(\mathbb{Z}_p[H]) \\ \pi_G \downarrow & & \downarrow \pi_H \\ K_i(\mathbb{Z}_p[G/N]) & \xrightarrow{\text{Nr}_{\mathbb{Z}_p[G/N]/\mathbb{Z}_p[H/N]}} & K_i(\mathbb{Z}_p[H/N]) \end{array}$$

where π_G and π_H are canonical homomorphisms induced by

$$\pi_G: \mathbb{Z}_p[G] \longrightarrow \mathbb{Z}_p[G/N] \quad \text{and} \quad \pi_H: \mathbb{Z}_p[H] \longrightarrow \mathbb{Z}_p[H/N].$$

When G is a pro-finite group, H is its open subgroup and N is a closed subgroup of H normal in G and H , the same statement holds for $\text{Nr}_{\Lambda(G)/\Lambda(H)}$ and $\text{Nr}_{\Lambda(G/N)/\Lambda(H/N)}$.

Proof. Let $\{u_1, \dots, u_r\}$ be a system of representatives of $H \backslash G$. Then it is clear that $\{\bar{u}_1, \dots, \bar{u}_r\}$ is that of $(H/N) \backslash (G/N)$ where $\bar{u}_i = \pi_G(u_i)$. Then we have

$$\begin{aligned} & \pi_H \circ \text{Nr}_{\mathbb{Z}_p[G]/\mathbb{Z}_p[H]} \\ &= [\mathbb{Z}_p[H/N] \otimes_{\mathbb{Z}_p[H]} (\mathbb{Z}_p[H] \mathbb{Z}_p[G]_{\mathbb{Z}_p[G]}) \otimes_{\mathbb{Z}_p[G]} -] \\ &= \left[\mathbb{Z}_p[H/N] \otimes_{\mathbb{Z}_p[H]} \left(\bigoplus_{i=1}^r \mathbb{Z}_p[H] u_i \right) \otimes_{\mathbb{Z}_p[G]} - \right] \\ &= \left[\left(\bigoplus_{i=1}^r \mathbb{Z}_p[H/N] \bar{u}_i \right) \otimes_{\mathbb{Z}_p[G]} - \right] \\ &= [(\mathbb{Z}_p[H/N] \mathbb{Z}_p[G/N]_{\mathbb{Z}_p[G/N]}) \otimes_{\mathbb{Z}_p[G/N]} (\mathbb{Z}_p[G/N]) \otimes_{\mathbb{Z}_p[G]} -] \\ &= \text{Nr}_{\mathbb{Z}_p[G/N]/\mathbb{Z}_p[H/N]} \circ \pi_G. \end{aligned}$$

The pro-finite version can be verified in the same manner. $\square$

1.3. The localization exact sequences. Let R be an associative ring and S be a multiplicatively closed subset of R containing 1. In 1968, Hyman Bass constructed “the K_1 - K_0 localization sequence” ([Bass1]):

$$K_1(R) \rightarrow K_1(S^{-1}R) \xrightarrow{\partial} K_0(R, S) \rightarrow K_0(R) \rightarrow K_0(S^{-1}R)$$

for central S (recall that S is *central* if S contained in the center of R).

In this subsection, we define Ore localizations (certain “good” localizations of non-commutative associative rings) and introduce the K_1 - K_0 localization exact sequence for a non-central Ore set S which is the generalization of Bass’ exact sequence for central S .

Definition 1.10 (Ore sets). Let R be an associative ring, and $S \subseteq R$ be a multiplicatively closed subset containing 1.

S is called a *left (resp. right) Ore set* if S satisfies following two conditions.

- (Ore-1) For every $r \in R$ and $s \in S$, $Sr \cap Rs \neq \emptyset$ (resp. $rS \cap sR \neq \emptyset$).
- (Ore-2) If $r \in R$ satisfies $rs = 0$ (resp. $sr = 0$) for a certain element $s \in S$, there exists $s' \in S$ such that $s'r = 0$ (resp. $rs' = 0$).

Example 1.11. Let R be an associative ring and S be a multiplicatively closed subset containing 1. Suppose that S is contained in the center of R . Then S is a left and right Ore set: for an arbitrary $r \in R$ and $s \in S$, there exist $r' \in R$ and $s' \in S$ such that $s'r = r's$ and $rs' = sr'$ since we may choose $s' = s$ and $r' = r$.

Proposition 1.12. Let R be an associative ring and S a left (resp. right) Ore set. Let $k \in \mathbb{N}$. Then the following property holds:

- (Ore-1') For arbitrary $s_1, \dots, s_k \in S$, there exist $r_1, \dots, r_k \in R$ which satisfy $r_1 s_1 = \dots = r_k s_k \in S$ (resp. $s_1 r_1 = \dots = s_k r_k \in S$).

Proof. Directly by (Ore-1) for $k = 2$. Then we can show (Ore-1') for $k \geq 3$ by induction. $\square$

Proposition-Definition 1.13 (Ore localization). Let R be an associative ring and S be a left (resp. right) Ore set of R .

- (1) (Existence of the Ore localization)

There exist a ring $[S^{-1}]R$ (resp. $R[S^{-1}]$) and a canonical ring homomorphism $\iota: R \rightarrow [S^{-1}]R$ (resp. $R[S^{-1}]$) which satisfy the following conditions:

- (loc-1) For every $s \in S$, $\iota(s)$ is invertible.
- (loc-2) Every element in $[S^{-1}]R$ (resp. $R[S^{-1}]$) can be described in the form $\iota(s)^{-1}\iota(r)$ (resp. $\iota(r)\iota(s)^{-1}$) for certain $r \in R$ and $s \in S$.
- (loc-3) For $r \in R$, $\iota(r) = 0$ if and only if there exists $s \in S$ such that $sr = 0$ (resp. $rs = 0$) in R .

- (2) (The universality property)

Suppose that a ring homomorphism $f: R \rightarrow \tilde{R}$ maps all elements of S to invertible elements of $\tilde{R}$. Then there exists a unique ring homomorphism ψ which makes the following diagram commute:

$$\begin{array}{ccc}
& [S^{-1}]R & (\text{resp. } R[S^{-1}]) \\
& \nearrow \iota & \downarrow \exists! \psi \\
R & \xrightarrow{f} & \widetilde{R}
\end{array}$$

We call $[S^{-1}]R$ (resp. $R[S^{-1}]$) the left (resp. right) Ore localization of R with respect to S .

For an arbitrary left (resp. right) R -module M , we define the left (resp. right) Ore localization of M with respect to S to be the module $[S^{-1}]M = [S^{-1}]R \otimes_R M$ (resp. $M[S^{-1}] = M \otimes_R R[S^{-1}]$).

Sketch of the proof. We only give the construction of the left Ore localization $[S^{-1}]R$. Set

$$[S^{-1}]R = S \times R / \sim$$

where $\sim$ is an equivalence relation defined by $(s, r) \sim (s', r')$ if and only if there exist $a, b \in R$ which satisfy $ar = br'$ and $as = bs' \in S$.

Then we may define the additive law and the multiplicative law as follows:

$$\begin{aligned}
(s, r) + (s', r') &= (t, ar + br') & \text{where } t = as = bs' \in S, \\
(s, r) \cdot (s', r') &= (ts, ar') & \text{where } tr = as', \quad t \in S.
\end{aligned}$$

For the well-definedness of $\sim$, $+$, $\cdot$, and for the universality property, we use the Ore conditions (Ore-1) and (Ore-2).

For details, see [Sten], Chapter II. $\square$

Corollary 1.14. *Let R be an associative ring. If a multiplicatively closed subset S satisfies both left and right Ore conditions, then the canonical isomorphism $[S^{-1}]R \xrightarrow{\sim} R[S^{-1}]$ exists.*

Proof. Directly from the universality property. $\square$

Proposition 1.15. *Let R be an associative ring and S a left (resp. right) Ore set. Then the left (resp. right) Ore localization $[S^{-1}]R$ (resp. $R[S^{-1}]$) is flat as a right (resp. left) R -module.*

In other words, the left (resp. right) Ore localization defines the exact functor from the category of left (resp. right) R -modules to that of left $[S^{-1}]R$ -modules (resp. right $R[S^{-1}]$ -modules).

Proof. See [Sten], Chapter II, Proposition 3.5. $\square$

Definition 1.16 (S -torsion modules). Let R be an associative ring and S be a left Ore set. We define $\mathfrak{M}_S(R)$ to be the category of finitely generated S -torsion left R -modules, that is, an object M of $\mathfrak{M}_S(R)$ is a finitely generated left R -module satisfying $[S^{-1}]M = 0$.

We may easily show that for an arbitrary object M of $\mathfrak{M}_S(R)$, there exists an element of S such that $sM = 0$ if S is central.

Bass constructed the following localization exact sequence for central S .

Definition 1.17. Let R be an associative ring and S its multiplicatively closed subset. Let $\mathfrak{H}_S(R)$ be the category of finitely generated S -torsion left R -modules with projective resolutions of finite length. Then we put $K_0(R, S) = K_0(\mathfrak{H}_S(R))$.

Remark 1.18. Note that we may identify $K_0(R, S)$ with the relative Grothendieck group associated to the canonical ring homomorphism $R \rightarrow [S^{-1}]R$. For the definition of relative Grothendieck groups, see [Bass1], Chapter IX, §1.

We can also identify this group with the Grothendieck group of the category of bounded complexes of finitely generated projective left R -modules whose cohomologies are of S -torsion.

Theorem 1.19 (Bass, the localization exact sequence for central S). *Let R be an associative ring with 1 and let S be its multiplicatively closed subset contained in the center of R . Suppose that for every element s of S , multiplication by s in R induces an injection $R \xrightarrow{s} R$.*

Then there exists an exact sequence of K -groups:

$$K_1(R) \rightarrow K_1(S^{-1}R) \xrightarrow{\partial} K_0(R, S) \rightarrow K_0(R) \rightarrow K_0(S^{-1}R).$$

Proof. See [Bass1], Chapter IX, Theorem (6.3). $\square$

Proposition 1.20. *If R has finite global dimension (in other words, if every finitely generated left R -module has finite projective dimension), we have*

$$K_i(R, S) = K_i(\mathfrak{M}_S(R)) \quad (i = 0, 1).$$

Proof. It is clear since $\mathfrak{H}_S(R) = \mathfrak{M}_S(R)$ in this case. $\square$

A. J. Berrick and M. E. Keating constructed the localization sequence for Ore localizations by generalizing the Bass' construction of the localization sequence for central S :

Theorem 1.21 (Berrick-Keating). *Let R be an associative ring with 1 and S a left Ore set. Suppose that no elements of S are zero divisors in R .*

Let $\mathfrak{H}_{S,1}(R)$ be the subcategory of $\mathfrak{M}_S(R)$ consisting of finitely presented S -torsion left R -modules with projective dimension 1. Then there exists an exact sequence of K -groups:

$$(1.2) \quad K_1(R) \rightarrow K_1([S^{-1}]R) \xrightarrow{\partial} K_0(\mathfrak{H}_{S,1}(R)) \rightarrow K_0(R) \rightarrow K_0([S^{-1}]R).$$

Proof. See [Ber-Keat]. $\square$

We remark that Daniel R. Grayson also constructed the exact sequence (1.2) by using Quillen's higher K -theory ([Grayson]).

In the following, we use the modified version of Berrick-Keating's localization sequence.

Theorem 1.22 (The localization exact sequence for Ore localization). *Let R be an associative ring with 1 and S a left Ore set. Suppose that no elements of S are zero divisors in R . Then there exists an exact sequence of K -groups:*

$$K_1(R) \rightarrow K_1([S^{-1}]R) \xrightarrow{\partial} K_0(R, S) \rightarrow K_0(R) \rightarrow K_0([S^{-1}]R).$$

Proof. It suffices to show that

$$(1.3) \quad K_0(\mathfrak{H}_S(R)) = K_0(\mathfrak{H}_{S,1}(R)).$$

First, we show that for every $M \in \text{Ob } \mathfrak{H}_S(R)$, M has a finite resolution by objects of $\mathfrak{H}_{S,1}(R)$. Let $\{m_1, \dots, m_k\}$ be elements of M which generate M , and d the projective dimension of M . Then we have the canonical surjection

$$\pi: \bigoplus_{i=1}^k Re_i \rightarrow M; e_i \mapsto m_i \quad (1 \leq i \leq k),$$

where $\{e_i\}_{1 \leq i \leq k}$ is the free basis. We set $L = \text{Ker}(\pi)$. Note that L is a left free module. Since M is a S -torsion module, there exists $s_i \in S$ which annihilates m_i for each i . Then we obtain an element $s \in S$ which annihilates all m_i by $(\text{Ore-1}')$. For this s , we have $\pi(\bigoplus_{i=1}^k Rse_i) = 0$, therefore $\bigoplus_{i=1}^k Rse_i$ is a left free R -submodule of L . Consider

$$0 \rightarrow L / \bigoplus_{i=1}^k Rse_i \rightarrow \bigoplus_{i=1}^k Re_i / \bigoplus_{i=1}^k Rse_i \rightarrow M \rightarrow 0,$$

then $\bigoplus_{i=1}^k Re_i / \bigoplus_{i=1}^k Rse_i$ is free and of S -torsion (Let $x = \sum_{i=1}^k r_i e_i$ be an arbitrary element of $\bigoplus_{i=1}^k Re_i / \bigoplus_{i=1}^k Rse_i$. For each i there exist $s'_i \in S$ and $r'_i \in R$ such that $s'_i r_i = r'_i s$. We obtain $r''_i \in R$ ($1 \leq i \leq k$) which satisfies $r''_1 s'_1 = \dots = r''_k s'_k \in S$ by $(\text{Ore-1}')$. Then the element $\tilde{s} = r''_1 s'_1 = \dots = r''_k s'_k$ annihilates x), therefore $\bigoplus_{i=1}^k Re_i / \bigoplus_{i=1}^k Rse_i$ is an object of $\mathfrak{H}_{S,1}(R)$. Note that the projective dimension of $L / \bigoplus_{i=1}^k Rse_i$ is $d-1$. Hence we can construct a resolution of M by objects of $\mathfrak{H}_{S,1}(R)$ by induction on the projective dimension d .

Then (1.3) reduces to Grothendieck's resolution theorem (See Theorem 1.23). Note that both $\mathfrak{H}_S(R)$ and $\mathfrak{H}_{S,1}(R)$ are admissible subcategories of $\mathfrak{M}(R)$ (that is, $\mathfrak{H}_S(R)$ and $\mathfrak{H}_{S,1}(R)$ are full additive subcategories of $\mathfrak{M}_S(R)$ which have at most sets of isomorphism classes of objects, and if $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is an arbitrary exact sequence in $\mathfrak{M}(R)$ for which M and M' are objects of $\mathfrak{H}_S(R)$ (resp. $\mathfrak{H}_{S,1}(R)$), M'' is also an object of $\mathfrak{H}_S(R)$ (resp. $\mathfrak{H}_{S,1}(R)$)). See [Bass2], Corollary (8.5). $\square$

Theorem 1.23 (Grothendieck's resolution theorem). *Let $\mathfrak{M}$ be an abelian category and $\mathfrak{P} \subseteq \mathfrak{P}'$ admissible subcategories of $\mathfrak{M}$. Assume an arbitrary object P' of $\mathfrak{P}'$ has a finite resolution by objects of $\mathfrak{P}$. Then the inclusion $\mathfrak{P} \subseteq \mathfrak{P}'$ induces an isomorphism $K_0(\mathfrak{P}) \cong K_0(\mathfrak{P}')$.*

Proof. See [Bass2], Theorem (7.1). $\square$

Remark 1.24. In our case, the canonical Ore set S for $\Lambda(G)$ (See §2) is essentially contained in the center $\Lambda(\Gamma)$ (See §6, Lemma 6.1), therefore Bass' localization sequence is sufficient for the proof of our main theorem.

1.4. Theory of the integral logarithm. Integral logarithmic homomorphisms were used by Robert Oliver and Laurence R. Taylor to study structure of Whitehead groups of group rings of finite groups ([Oliver], [Oli-Tay]). We use these homomorphisms to translate “the additive theta map” into “the (multiplicative) theta map” (See §5). Ritter and Weiss also used them to formulate their “equivariant Iwasawa theory” ([R-W1]).

Let R be a complete discrete valuation ring with mixed characteristics $(0, p)$, K its fractional field, π a uniformizer of R and k the residue field of R . Let $\mathfrak{A}$ be an “ R -order,” that is, an R -algebra which is free as a left R -module (though Oliver and Taylor treated only $\mathbb{Z}_p$ -orders in [Oliver] and [Oli-Tay], we need integral logarithms for the $\widehat{\Lambda(\Gamma)_{(p)}}$ -order $\widehat{\Lambda(\Gamma)_{(p)}}[G^f]$ to construct

the localized version of the theta map (see § 6), therefore we introduce here the generalized version of integral logarithms for general R -orders).

Lemma 1.25 ([Oliver], Lemma 2.7 for $\mathbb{Z}_p$ -orders). *Let R , K , π , k and $\mathfrak{A}$ be as above, and let J be the Jacobson radical of $\mathfrak{A}$. Then for every two-sided ideal $\mathfrak{a} \subseteq J$, $1 + \mathfrak{a}$ is a multiplicative group. Moreover, for an arbitrary element $x \in \mathfrak{a}$, the power series*

$$(1.4) \quad \log(1+x) = \sum_{i=1}^{\infty} (-1)^{i-1} \frac{x^i}{i}$$

converges p -adically in $\mathfrak{a}_{\mathbb{Q}} = \mathbb{Q} \otimes_{\mathbb{Z}} \mathfrak{a}$, and satisfies

$$(1.5) \quad \log((1+x)(1+y)) \equiv \log(1+x) + \log(1+y) \pmod{[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}]}$$

for every $x, y \in \mathfrak{a}$ where $\mathfrak{A}_{\mathbb{Q}} = \mathbb{Q} \otimes_{\mathbb{Z}} \mathfrak{A}$.

In particular, $\log$ induces a homomorphism of groups

$$\log: 1 + \mathfrak{a} \longrightarrow \mathfrak{a}_{\mathbb{Q}}/[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}].$$

Here for arbitrary two-sided ideals $\mathfrak{a}$ and $\mathfrak{b}$ of R , $[\mathfrak{a}, \mathfrak{b}]$ denotes the two-sided ideal generated by $[a, b] = ab - ba$ where $a \in \mathfrak{a}$ and $b \in \mathfrak{b}$.

Proof. Since $\mathfrak{A}/\mathfrak{A}\pi$ is finite over $k = R/R\pi$, it is a left Artinian k -algebra. Note that $J/\mathfrak{A}\pi$ is the Jacobson radical of $\mathfrak{A}/\mathfrak{A}\pi$, and by commutative ring theory, the Jacobson radical of an arbitrary (left) Artinian ring is nilpotent. Hence for an arbitrary $x \in \mathfrak{a} \subseteq J$, there exists a certain natural number m such that

$$(1.6) \quad x^m \in \mathfrak{A}\pi.$$

Let e be the absolute ramification index of K . Then we have $x^n \in \mathfrak{A}p^{[n/em]}$ for each $n \geq em$ where $[r]$ denotes the greatest integer not greater than r for every $r \in \mathbb{R}$. $\mathfrak{A}p^{[n/em]}$ converges to zero as $n \rightarrow \infty$, therefore we obtain

$$\lim_{n \rightarrow \infty} |x^n|_p = 0,$$

which implies the power series

$$(1+x)^{-1} = \sum_{i=0}^{\infty} (-1)^i x^i$$

converges in $\mathfrak{A}$. Since $\mathfrak{a}$ is closed in p -adic topology, we may conclude that $(1+x)^{-1} \in 1 + \mathfrak{a}$. It is clear that $1 + \mathfrak{a}$ is closed under multiplication (note that $\mathfrak{a}$ is a two-sided ideal), therefore $1 + \mathfrak{a}$ is a multiplicative group.

Now we also have $x^n/n \in \mathfrak{A} \cdot (p^{[n/em]}/n)$ for each $n \geq em$ by (1.6), therefore we obtain

$$(1.7) \quad \lim_{n \rightarrow \infty} \left| \frac{x^n}{n} \right|_p = 0,$$

which implies that the power series (1.4) converges in $\mathfrak{A}_{\mathbb{Q}}$. Then we can show that (1.4) converges in $\mathfrak{a}_{\mathbb{Q}}$ by the same argument as above.

We may also show the equation (1.5) by direct calculation, but this calculation is quite complicated because of the non-commutativity of $\mathfrak{A}$. See [Oliver], Chapter 2, Lemma 2.7. $\square$

Remark 1.26. Note that though Oliver constructed $\log$ only for $\mathbb{Z}_p$ -orders in [Oliver], Lemma 2.7 of Chapter 2, his proof does not use peculiarities of $\mathbb{Z}_p$ (especially the finiteness of the residue field of R). So we can apply his proof to the case of general R -orders.

Proposition 1.27 ([Oliver], Theorem 2.8 for $\mathbb{Z}_p$ -orders). *Let $\mathfrak{A}$ and $\mathfrak{a}$ be as above. Then the logarithmic homomorphism of Lemma 1.25 induces the following homomorphism of abelian groups:*

$$\log_{\mathfrak{a}}: K_1(\mathfrak{A}, \mathfrak{a}) \longrightarrow \mathfrak{a}_{\mathbb{Q}}/[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}].$$

Sketch of the proof. For every $n \geq 1$ and for an ideal $M_n(\mathfrak{a}) \subseteq M_n(\mathfrak{A})$, we have a logarithmic homomorphism (Lemma 1.25)

$$\begin{aligned} \log^{(n)}: \mathrm{GL}_n(\mathfrak{A}, \mathfrak{a}) = I_n + M_n(\mathfrak{a}) &\xrightarrow{\log} M_n(\mathfrak{a}_{\mathbb{Q}})/[M_n(\mathfrak{A}_{\mathbb{Q}}), M_n(\mathfrak{a}_{\mathbb{Q}})] \\ &\xrightarrow{\text{trace}} \mathfrak{a}_{\mathbb{Q}}/[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}] \end{aligned}$$

where I_n is the unit matrix of $\mathrm{GL}_n(\mathfrak{A})$.

For every $X \in \mathrm{GL}_n(\mathfrak{A})$ and $A \in \mathrm{GL}_n(\mathfrak{A}, \mathfrak{a}) = I_n + M_n(\mathfrak{a})$, we have

$$\begin{aligned} \log^{(n)}([X, A]) &= \log^{(n)}(XAX^{-1}) - \log^{(n)}(A) \quad (\text{by (1.5)}) \\ &= \mathrm{Tr}(X \log(A)X^{-1}) - \mathrm{Tr}(\log(A)) \\ &= 0. \end{aligned}$$

Hence $\log^{(n)}$ induces

$$\log^{(n)}: \mathrm{GL}_n(\mathfrak{A})/[\mathrm{GL}_n(\mathfrak{A}), \mathrm{GL}_n(\mathfrak{A}, \mathfrak{a})] \longrightarrow \mathfrak{a}_{\mathbb{Q}}/[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}],$$

and by taking the projective limit, we obtain

$$\log_{\mathfrak{a}}: K_1(\mathfrak{A}, \mathfrak{a}) \longrightarrow \mathfrak{a}_{\mathbb{Q}}/[\mathfrak{A}_{\mathbb{Q}}, \mathfrak{a}_{\mathbb{Q}}],$$

using Whitehead's lemma (Proposition 1.5).

(See [Oliver], Theorem 2.8.) □

Remark 1.28. Let R be the integer ring of a finite extension of $\mathbb{Q}_p$, and take $\mathfrak{a}$ to be the Jacobson radical J of $\mathfrak{A} = R[G]$ where G is a finite group. Then we may extend the domain of the logarithmic homomorphism obtained in Proposition 1.27 to $K_1(R[G])$ uniquely.

The following exact sequence of K -groups is well known:

$$K_2(R[G]/J) \rightarrow K_1(R[G], J) \rightarrow K_1(R[G]) \rightarrow K_1(R[G]/J) \rightarrow 1$$

(See [Milnor] Lemma 4.1 and Theorem 6.2). Note that the homomorphism $K_1(R[G]) \rightarrow K_1(R[G]/J)$ is surjective by Proposition 1.6. Since $R[G]/J$ is a finite semi-simple algebra with p -power order, $K_2(R[G]/J) = 1$ and $p \nmid a = \#K_1(R[G]/J)$ ([Oliver], Theorem 1.16). Therefore, we may regard $K_1(R[G], J)$ as a subgroup of $K_1(R[G])$, and may extend the domain of $\log_J$ to $K_1(R[G])$ by setting

$$\log_{R[G]} \phi = \frac{1}{a} \log_J \phi^a$$

for every $\phi \in K_1(R[G])$ (note that $\phi^a \in K_1(R[G], J)$). The uniqueness of the extension is trivial from this construction.

Now we define *integral logarithmic homomorphisms*. Let R be a complete discrete valuation ring which is absolutely unramified, and let G be a finite p -group. For simplicity, we assume that $p \neq 2$. We consider a group ring $R[G]$. Let J be the Jacobson radical of $R[G]$. Let

$$R[\text{Conj}(G)] = R[G]/[R[G], R[G]] \quad (\text{resp. } K[\text{Conj}(G)] = K[G]/[K[G], K[G]])$$

be the free R -module (resp. the K -vector space) with basis $\text{Conj}(G)$. In the following, we fix the Frobenius automorphism $\tilde{\varphi}: K \rightarrow K$ when $k = R/pR$ is not perfect. Then we have

$$(1.8) \quad \tilde{\varphi}(r) \equiv r^p \pmod{pR}$$

for every $r \in R$.²

Let $\varphi: K[\text{Conj}(G)] \rightarrow K[G]$ be the homomorphism defined by

$$\varphi\left(\sum_g k_g [g]\right) = \sum_g \tilde{\varphi}(k_g) [g^p] \quad (k_g \in K, [g] \in \text{Conj}(G)).$$

Proposition-Definition 1.29 (The integral logarithms). *Set*

$$\Gamma_{G,J}(u) = \log_J(u) - \frac{1}{p}\varphi(\log_J(u)) \in K[\text{Conj}(G)], \quad u \in K_1(R[G], J).$$

Then $\Gamma_{G,J}$ induces a homomorphism

$$\Gamma_{G,J}: K_1(R[G], J) \rightarrow R[\text{Conj}(G)],$$

which we call *the integral logarithmic homomorphism for $R[G]$* .

When K is a finite unramified extension of $\mathbb{Q}_p$ and R is its integer ring, then

$$\Gamma_G(u) = \log_{R[G]}(u) - \frac{1}{p}\varphi(\log_{R[G]}(u)) \in K[\text{Conj}(G)], \quad u \in K_1(R[G])$$

also induces a homomorphism

$$\Gamma_G: K_1(R[G]) \rightarrow R[\text{Conj}(G)].$$

Sketch of the proof. Take an arbitrary $x \in J$. Since

$$\begin{aligned} \Gamma_{G,J}(1-x) &= -\left(x + \frac{x^2}{2} + \cdots\right) + \frac{1}{p}\varphi\left(x + \frac{x^2}{2} \cdots\right) \\ &\equiv -\sum_{i=1}^{\infty} \frac{1}{pi} (x^{pi} - \varphi(x^i)) \pmod{R[\text{Conj}(G)]}, \end{aligned}$$

it is sufficient to show that $pi|(x^{pi} - \varphi(x^i))$ for every $i \geq 1$, or $p^n|(x^{p^n} - \varphi(x^{p^{n-1}}))$ for every $n \geq 1$ (note that all primes other than p are invertible in R).

To do this, we should check each term of the expansion of x^{p^n} and $\varphi(x^{p^{n-1}})$ carefully. See [Oliver], Theorem 6.2 for details.

The second part follows immediately by the first part and the uniqueness of the extension of $\log$ (Remark 1.28). $\square$

²If R is the integer ring of a finite unramified extension of $\mathbb{Q}_p$, $\tilde{\varphi}$ is the ordinary Frobenius endomorphism (determined uniquely up to the inertia subgroup). If R is the p -adic completion of $\Lambda(\Gamma)_{(p)}$ (we use this ring in §6), we may choose $\tilde{\varphi}$ as the endomorphism induced by $t \mapsto t^p$.

Now assume that R is the p -adic integer ring $\mathbb{Z}_p$ and G is a finite p -group. In this case we can derive further information about the kernel and image of the integral logarithms.

Theorem 1.30. *Let G be a finite p -group and $K_1(\mathbb{Z}_p[G])_{\text{tors}}$ the torsion part of $K_1(\mathbb{Z}_p[G])$. Then there exists an exact sequence*

$$1 \rightarrow K_1(\mathbb{Z}_p[G])/K_1(\mathbb{Z}_p[G])_{\text{tors}} \xrightarrow{\Gamma_G} \mathbb{Z}_p[\text{Conj}(G)] \xrightarrow{\omega_G} G^{\text{ab}} \rightarrow 1.$$

Here ω_G is defined by

$$\omega_G \left(\sum_g a_g [g] \right) = \prod_g \bar{g}^{a_g} \quad (a_g \in \mathbb{Z}_p, [g] \in \text{Conj}(G))$$

and G^{ab} is the abelization of G . We denote by $\bar{g}$ the image of $[g]$ in G^{ab} .

Proof. See [Oliver], Theorem 6.6. $\square$

For the torsion part of $K_1(\mathbb{Z}_p[G])$, the following properties are known.

Definition 1.31 ($SK_1(\mathbb{Z}_p[G])$). Let G be a finite group. Then we put

$$SK_1(\mathbb{Z}_p[G]) = \text{Ker} (K_1(\mathbb{Z}_p[G]) \rightarrow K_1(\mathbb{Q}_p[G])).$$

Remark 1.32. We may define the SK_1 -group for an arbitrary associative ring by Whitehead construction, but we omit this since we only use SK_1 -groups for group rings of $\mathbb{Z}_p[G]$ -type.

Proposition 1.33. *Let G be a finite group.*

- (1) $K_1(\mathbb{Z}_p[G])_{\text{tors}} \cong \mu_{p-1} \times G^{\text{ab}} \times SK_1(\mathbb{Z}_p[G])$.
- (2) $SK_1(\mathbb{Z}_p[G])$ is finite.
- (3) If G is commutative, $SK_1(\mathbb{Z}_p[G]) = 1$.

Proof. (1) See [Oliver], Theorem 7.4.

(2) See [Wall], Theorem 2.5.

(3) This follows from the definition of SK_1 -groups: since $\mathbb{Z}_p[G]$ and $\mathbb{Q}_p[G]$ are local and commutative, we have $K_1(\mathbb{Z}_p[G]) \cong \mathbb{Z}_p[G]^\times$ and $K_1(\mathbb{Q}_p[G]) \cong \mathbb{Q}_p[G]^\times$ by Proposition 1.7 (1), then $\mathbb{Z}_p[G]^\times \rightarrow \mathbb{Q}_p[G]^\times$ is obviously injective. $\square$

2. BASIC RESULTS OF NON-COMMUTATIVE IWASAWA THEORY

In this section, we review basic results of [CFKSV].

Let G be a compact p -adic Lie group containing a normal closed subgroup H which satisfies $G/H \cong \mathbb{Z}_p$, and let $\Lambda(G)$ be its Iwasawa algebra. In the non-commutative Iwasawa theory of [CFKSV], characteristic elements (the ‘‘arithmetic’’ p -adic zeta functions) are defined as elements of Whitehead groups of certain Ore localization of $\Lambda(G)$ by using the localization exact sequence. We first define the *canonical Ore set* S for the group G introduced in [CFKSV] §2. Then for every element $[\mathfrak{C}]$ of the relative K_0 -group $K_0(\Lambda(G), \Lambda(G)_S)$ (in the following, we regard $K_0(\Lambda(G), \Lambda(G)_S)$ as the Grothendieck group of the category of bounded complexes of finitely generated projective left $\Lambda(G)$ -modules whose cohomologies are of S -torsion), we define a *characteristic element of* $[\mathfrak{C}]$ as an element of $K_1(\Lambda(G)_S)$. Next,

we characterize the p -adic zeta function (the “analytic” element) as an element of $K_1(\Lambda(G)_S)$ interpolating special values of Artin L -functions, and formulate the non-commutative Iwasawa main conjecture.

2.1. The canonical Ore set and characteristic elements. Let G be a compact p -adic Lie group and

$$\Lambda(G) = \varprojlim_{U: \text{open normal}} \mathbb{Z}_p[G/U]$$

be its Iwasawa algebra. Suppose that there exists a normal closed subgroup H of G which satisfies $G/H \cong \Gamma$, where Γ is a commutative p -adic Lie group isomorphic to $\mathbb{Z}_p$ (See §0.3). In the following, we fix such a subgroup H .

Proposition-Definition 2.1 (The canonical Ore set). *Let G and H be as above. Then*

$$S = \{f \in \Lambda(G) \mid \Lambda(G)/\Lambda(G)f \text{ is finitely generated as a left } \Lambda(H)\text{-module}\}$$

is a left and right Ore set (See Definition 1.10).

We call this S the canonical Ore set for the group G .³

Since S is a left and right Ore set, the left localization and the right localization of $\Lambda(G)$ with respect to S are canonically isomorphic to each other by Corollary 1.14. Therefore we may identify these two localizations. Because of this reason, we denote by $\Lambda(G)_S$ the left (or right) localization of $\Lambda(G)$ with respect to S .

The canonical Ore sets and their properties are discussed well in [CFKSV], §2. Here we use the following two properties:

Proposition 2.2. *Let G be a compact p -adic Lie group and S the canonical Ore set for it.*

- (1) *The localized Iwasawa algebra $\Lambda(G)_S$ is semi-local.*
- (2) *The elements of S are non-zero divisors in $\Lambda(G)$.*

Proof. (1) See [CFKSV], Proposition 4.2.

(2) See [CFKSV], Theorem 2.4. □

Note that we have the canonical surjection $\Lambda(G)_S^\times \rightarrow K_1(\Lambda(G)_S)$ by Proposition 1.7 (1) and Proposition 2.2 (1). Also note that by Proposition 2.2 (2), we may consider the localization sequence for the Ore localization $\Lambda(G) \rightarrow \Lambda(G)_S$ (Theorem 1.22).

Remark 2.3. Assume that G has no p -torsion elements. In this case we often use

$$S^* = \bigcup_{n \geq 0} p^n S$$

in place of S , as is remarked in [CFKSV]. We may identify $K_0(\Lambda(G), \Lambda(G)_{S^*})$ with $K_0(\mathfrak{M}_{S^*}(\Lambda(G)))$ where $\mathfrak{M}_{S^*}(\Lambda(G))$ is the category of finitely generated left S^* -torsion $\Lambda(G)$ -modules.

³Since the definition of S depends on the subgroup H , we should call S the canonical Ore set for (G, H) . But by abuse of notation, we often call S the canonical Ore set for G when the subgroup H is fixed.

But in our case (See § 3), G has many p -torsion elements. Therefore we have to treat derived categories of complexes of finitely generated modules.

Now let us consider the localization exact sequence (Theorem 1.22) for $\Lambda(G) \rightarrow \Lambda(G)_S$:

$$(2.1) \quad K_1(\Lambda(G)) \rightarrow K_1(\Lambda(G)_S) \xrightarrow{\partial} K_0(\Lambda(G), \Lambda(G)_S) \rightarrow K_0(\Lambda(G)) \rightarrow K_0(\Lambda(G)_S).$$

Proposition 2.4. *The connecting homomorphism ∂ in (2.1) is surjective.*

Therefore, for an arbitrary element $[\mathfrak{C}] \in K_0(\Lambda(G), \Lambda(G)_S)$, there exists an element $f \in K_1(\Lambda(G)_S)$ which satisfies $\partial(f) = -[\mathfrak{C}]$.

Definition 2.5 (Characteristic element). Let $[\mathfrak{C}] \in K_0(\Lambda(G), \Lambda(G)_S)$. We call an element f of $K_1(\Lambda(G)_S)$ a *characteristic element* of $[\mathfrak{C}]$ if f satisfies $\partial(f) = -[\mathfrak{C}]$.

Remark 2.6. By the localization exact sequence, characteristic elements of $[\mathfrak{C}]$ are determined up to multiplication by elements of $K_1(\Lambda(G))$.

Proof of Proposition 2.4. We may prove this proposition by the almost same argument as the proof of Proposition 3.4 in [CFKSV].

Though we assume that G has no p -torsion elements in the proof of Proposition 3.4 in [CFKSV], this assumption is used only to avoid treating complexes directly (this point is also remarked in [CFKSV]). For each complex $\mathfrak{C} = \mathfrak{C}^\bullet$, its canonical image in $K_0(\Lambda(G))$ is $\sum_{i \in \mathbb{Z}} (-1)^i [\mathfrak{C}^i]$. By using this fact

and translating $\lambda, \tau, \varepsilon$ in [CFKSV] appropriately, we may apply their proof of Proposition 3.4 in [CFKSV] to the case where G has p -torsion elements. $\square$

Example 2.7. Let us consider the classical Iwasawa $\mathbb{Z}_p$ -extension case: $G = \Gamma, H = \{1\}$ and $\Lambda(\Gamma) \cong \mathbb{Z}_p[[T]]$; $t \mapsto 1 + T$.

In this case, by p -adic Weierstrass' preparation theorem, we have

$$f(T) = u(T)p^n g(T)$$

for an arbitrary non-zero element $f(T) \in \mathbb{Z}_p[[T]]$, where $u(T) \in \mathbb{Z}_p[[T]]^\times$, $n \in \mathbb{Z}_{\geq 0}$, and $g \in \mathbb{Z}_p[T]$ is a distinguished polynomial. Hence $\Lambda(\Gamma)/\Lambda(\Gamma)f$ is finitely generated over $\mathbb{Z}_p$ if and only if $p \nmid f$ (or equivalently $n = 0$). Therefore the canonical Ore set S is $\Lambda(\Gamma) \setminus p\Lambda(\Gamma)$.

Since Γ has no p -torsion elements, we may consider the Ore set S^* as in Remark 2.3. It is easy to see that in this case S^* is the subset of $\Lambda(\Gamma)$ consisting of all non-zero elements. Therefore $\Lambda(\Gamma)_{S^*}$ coincides with the fractional field $\text{Frac}(\Lambda(\Gamma))$.

Furthermore, since $\Lambda(\Gamma)$ is local and commutative, we have

$$K_1(\Lambda(\Gamma)) = \Lambda(\Gamma)^\times \quad \text{and} \quad K_1(\Lambda(\Gamma)_{S^*}) = \text{Frac}(\Lambda(\Gamma))^\times.$$

Hence the localization exact sequence is described as follows:

$$\Lambda(\Gamma)^\times \rightarrow \text{Frac}(\Lambda(\Gamma))^\times \xrightarrow{\partial} K_0(\mathfrak{M}_{\text{tor}}(\Lambda(\Gamma))) \rightarrow 0,$$

where $\mathfrak{M}_{\text{tor}}(\Lambda(\Gamma))$ is the category of all finitely generated torsion $\Lambda(\Gamma)$ -modules. The connecting homomorphism ∂ is characterized by $\partial(f) = [\Lambda(\Gamma)/\Lambda(\Gamma)f]$ for $f \in \Lambda(\Gamma) \setminus \{0\}$ in this (abelian) case.

Let M be an arbitrary finitely generated torsion $\Lambda(\Gamma)$ -module. Then by the famous structure theorem of finitely generated torsion $\Lambda(\Gamma)$ -modules, there exist $f_i \in \Lambda(\Gamma) \setminus \{0\}$ ($1 \leq i \leq N$) and satisfy

$$M \sim \bigoplus_{i=1}^N \Lambda(\Gamma)/\Lambda(\Gamma)f_i \quad (\text{pseudo-isomorphic})$$

where the image of each f_i in $\mathbb{Z}_p[[T]]$ is a non-invertible polynomial. Since the image of every pseudo-null $\Lambda(\Gamma)$ -module in $K_0(\mathfrak{M}_{\text{tor}}(\Lambda(\Gamma)))$ vanishes (See [Sch-Ven]), we have

$$[M] = \sum_{i=1}^N [\Lambda(\Gamma)/\Lambda(\Gamma)f_i].$$

Hence, by the explicit description of ∂ , we have

$$\partial(f_M) = [M]$$

where $f_M = \prod_{i=1}^N f_i$ (*the characteristic polynomial of M*). f_M is determined up to multiplication by an element of $\Lambda(\Gamma)^\times$.

The calculation above shows that the characteristic elements in [CFKSV] are generalized notion of the classical characteristic polynomials.

2.2. The non-commutative Iwasawa main conjecture for totally real fields. Now let us review the formulation of the non-commutative Iwasawa main conjecture in the sense of [CFKSV].

Fix a prime number $p \neq 2$. Let F be a totally real number field and F^∞/F a Galois extension of infinite degree satisfying the following conditions:⁴

- (1) The Galois group $G = \text{Gal}(F^\infty/F)$ is a compact p -adic Lie group.
- (2) Only finitely many primes of F ramify in F^∞ .
- (3) F^∞ is totally real and contains the cyclotomic $\mathbb{Z}_p$ -extension F^{cyc} of F .

Fix a finite set Σ of primes of F containing all primes which ramify in F^∞ .

Definition 2.8. Under the conditions above, we define the complex C by

$$\begin{aligned} C &= C_{F^\infty/F} \\ &= R\text{Hom}(R\Gamma_{\text{ét}}(\text{Spec } \mathcal{O}_{F^\infty}[1/\Sigma], \mathbb{Q}_p/\mathbb{Z}_p), \mathbb{Q}_p/\mathbb{Z}_p). \end{aligned}$$

Here $\Gamma_{\text{ét}}$ is the global section functor for étale topology.

Note that $H^0(C) = \mathbb{Z}_p$, $H^{-1}(C) = \text{Gal}(M_\Sigma/F^\infty)$ where M_Σ is the maximal abelian pro- p extension of F^∞ unramified outside Σ , and $H^n(C) = 0$ for $n \neq 0, -1$. We denote $\text{Gal}(M_\Sigma/F^\infty)$ by $X_\Sigma(F^\infty/F)$.

Now set $H = \text{Gal}(F^\infty/F^{\text{cyc}})$ and $\Gamma = \text{Gal}(F^{\text{cyc}}/F) \cong \mathbb{Z}_p$. Then if $[C]$ is an element of $K_0(\Lambda(G), \Lambda(G)_S)$, we can apply the results of §2.1 to $[C]$ and obtain a *characteristic element for F^∞/F* as a characteristic element of $[C]$.

Conjecture 2.9. $[C]$ is always an element of $K_0(\Lambda(G), \Lambda(G)_S)$. In other words, $X_\Sigma(F^\infty/F)$ is of S -torsion.

⁴Since the Galois extension we will consider has p -torsion elements, we need to weaken the conditions in [CFKSV]. See [Kato1] §2 and [Fu-Ka] §4.3.

Proposition 2.10. *Let $G' \subseteq G$ be a pro- p subgroup of G and let F' be the maximal intermediate field of F^∞/F fixed by G' , then the followings are equivalent:*

- (1) $X_\Sigma(F^\infty/F)$ is of S -torsion.
- (2) $\mu(F'^{\text{cyc}}/F') = 0$ where μ is the μ -invariant.

In particular, if the following condition $(*)$ is satisfied, $X_\Sigma(F^\infty/F)$ is of S -torsion:

- $(*)$ *There exists a finite subextension F' of F^∞ such that $\text{Gal}(F^\infty/F')$ is pro- p and $\mu(F'^{\text{cyc}}/F') = 0$.*

Proof. This proposition is a variant of [Ha-Sh], Lemma 3.4. $\square$

For the μ -invariants of cyclotomic $\mathbb{Z}_p$ -extensions, Kenkichi Iwasawa conjectured:

Conjecture 2.11 (Iwasawa's $\mu = 0$ conjecture). *For every number field K , $\mu(K^{\text{cyc}}/K) = 0$.*

Corollary 2.12. *Assume that Iwasawa's $\mu = 0$ conjecture is true, then $X_\Sigma(F^\infty/F)$ is always of S -torsion.*

Corollary 2.13. *Let $K/\mathbb{Q}$ be a finite abelian extension. Then $X_\Sigma(K^\infty/K)$ is of S -torsion.*

Proof. $\mu(K^{\text{cyc}}/K) = 0$ by Ferrero-Washington's theorem ([FW]). $\square$

In the following, we always assume the condition $(*)$ in Proposition 2.10.

Now we define the “ p -adic zeta function” as an element of $K_1(\Lambda(G)_S)$. Let

$$\rho: G \longrightarrow \text{GL}_d(\overline{\mathbb{Q}}) \rightarrow \text{GL}_d(\overline{\mathbb{Q}}_p)$$

be an arbitrary Artin representation (that is, $\rho(G) \subseteq \text{GL}_d(\overline{\mathbb{Q}})$ is a finite subgroup). Then there exists a finite extension E of $\mathbb{Q}_p$ such that $\text{GL}_d(E)$ contains the image of ρ , and ρ induces a ring homomorphism

$$\rho: \Lambda(G) \longrightarrow \text{M}_d(E)$$

This also induces a homomorphism of K -groups

$$\text{ev}_\rho: K_1(\Lambda(G)) \longrightarrow K_1(\text{M}_d(E)) \xrightarrow{\cong} K_1(E) \cong E^\times$$

where the isomorphism $K_1(\text{M}_d(E)) \xrightarrow{\cong} K_1(E)$ is given by the Morita equivalence between $\text{M}_d(E)$ and E . Composing this with the natural inclusion $E^\times \rightarrow \overline{\mathbb{Q}}_p^\times$, we obtain the map

$$\text{ev}_\rho: K_1(\Lambda(G)) \longrightarrow \overline{\mathbb{Q}}_p^\times.$$

As is discussed in [CFKSV] §2, this map can be extended to

$$\text{ev}_\rho: K_1(\Lambda(G)_S) \longrightarrow \overline{\mathbb{Q}}_p \cup \{\infty\}.$$

We call ev_ρ the *evaluation map at ρ* . We denote $\text{ev}_\rho(f)$ by $f(\rho)$ for every $f \in K_1(\Lambda(G)_S)$.

Let

$$\kappa: \text{Gal}(F(\mu_{p^\infty})/F) \longrightarrow \mathbb{Z}_p^\times$$

be the p -adic cyclotomic character. Then for every positive integer r divisible by $p-1$, κ^r factors $\Gamma = \text{Gal}(F^{\text{cyc}}/F)$. Therefore we may extend the domain of κ^r to G by

$$G = \text{Gal}(F^\infty/F) \rightarrow \text{Gal}(F^{\text{cyc}}/F) \xrightarrow{\kappa^r} \mathbb{Z}_p^\times$$

where the first map is the canonical surjection.

Definition 2.14 (p -adic zeta function). If $\xi_{F^\infty/F} \in K_1(\Lambda(G)_S)$ satisfies

$$(2.2) \quad \xi_{F^\infty/F}(\rho \otimes \kappa^r) = L_\Sigma(1-r; F^\infty/F, \rho)$$

for every positive integer r divisible by $p-1$ and for an arbitrary Artin representation ρ of G , we call $\xi_{F^\infty/F}$ the p -adic zeta function for F^∞/F .

Here $L_\Sigma(s; F^\infty/F, \rho)$ is the complex Artin L -function of ρ in which the Euler factors at Σ are removed.

Conjecture 2.15. Let F^∞/F be as above.

- (1) (*The existence and the uniqueness of the p -adic zeta function*)
The p -adic zeta function $\xi_{F^\infty/F}$ for F^∞/F exists uniquely.
- (2) (*The non-commutative Iwasawa main conjecture*)
The p -adic zeta function $\xi_{F^\infty/F}$ satisfies $\partial(\xi_{F^\infty/F}) = -[C_{F^\infty/F}]$.

Remark 2.16 (The abelian case). Let $G = \text{Gal}(F^\infty/F)$ be an abelian p -adic Lie group. In this case, Coates observed that if certain congruences among the special values of the partial zeta functions were proven, we could construct the p -adic L -function for F^∞/F (See [Coates], Hypotheses (H_n) and (C_0)). These congruences were proven by Deligne and Ribet using the deep result about Hilbert-Blumenthal modular varieties ([De-Ri]).

Using the Deligne-Ribet's congruences, Serre constructed the element $\xi_{F^\infty/F}$ of $\text{Frac}(\Lambda(G))$ which satisfied the following two properties (Serre's p -adic zeta pseudomeasure [Serre2] for F^∞/F , also see §7).

- (1) For an arbitrary element g of G , $(1-g)\xi_{F^\infty/F}$ is contained in $\Lambda(G)$.
- (2) $\xi_{F^\infty/F}$ satisfies the interpolation property (2.2).

Remark 2.17. The non-commutative Iwasawa main conjecture which we introduced here was established first by Coates, Fukaya, Kato, Sujatha and Venjakob for elliptic curves without complex multiplication (the GL_2 -conjecture) in [CFKSV]. For more precise statements, see [CFKSV].

In [Fu-Ka], Fukaya and Kato established the main conjecture for general cases and showed the compatibility of the main conjecture with the equivariant Tamagawa number conjecture.

3. THE MAIN THEOREM AND BURNS' TECHNIQUE

3.1. The main theorem. Fix a prime number p . We consider the one dimensional p -adic Lie group $G = G^f \times \Gamma$ where

$$G^f = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (\text{finite part of } G)$$

and Γ is a commutative p -adic Lie group isomorphic to $\mathbb{Z}_p$ (See §0.3).

In the following, we fix generators of G^f and denote them by

$$\begin{aligned}\alpha &= \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \beta &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ \gamma &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \delta &= \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ \varepsilon &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \zeta &= \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.\end{aligned}$$

Then the center of G^f is $\langle \zeta \rangle$ and there are four non-trivial fundamental relations of G^f :

$$\begin{aligned}[\alpha, \beta] &= \delta, & [\beta, \gamma] &= \varepsilon, \\ [\alpha, \varepsilon] &= \zeta, & [\delta, \gamma] &= \zeta,\end{aligned}$$

where $[x, y] = xyx^{-1}y^{-1}$ is the commutator of x and y . Other commutators among $\alpha, \beta, \dots, \zeta$ are 1. We always denote the indices of $\alpha, \beta, \dots, \zeta$ by $a, b, \dots, f$. Note that G satisfies the conditions in §2 for $H = G^f$. We also assume that $p \neq 2, 3$. Under this assumption, the exponent of the group G^f is p .

Let F be a totally real number field and F^∞ a Galois extension of F satisfying $\text{Gal}(F^\infty/F) \cong G$ and the condition $(*)$ in Proposition 2.10, that is, there exists an intermediate field $F^\infty/F'/F$ such that F'/F is finite and the μ -invariant $\mu((F')^{\text{cyc}}/F')$ equals to zero.

Theorem 3.1. *Under the notation and the assumptions as above, the p -adic zeta function $\xi_{F^\infty/F}$ for F^∞/F exists and the main conjecture (Conjecture 2.15 (2)) is true.*

Remark 3.2. In this paper we do not discuss the uniqueness of $\xi_{F^\infty/F}$. Also see Remark 5.15.

3.2. Burns' technique. In this subsection, let F^∞/F be a general p -adic Lie extension of a totally real field F satisfying the conditions in §2.2. Put $G = \text{Gal}(F^\infty/F)$.

Let $\mathfrak{F}$ be a family of pairs (U, V) where U is an open subgroup of G and V is an open subgroup of H such that V is normal in U and U/V is commutative.

For $\mathfrak{F}$, we assume the following hypothesis:

- (b) For an arbitrary Artin representation ρ of G , ρ is a $\mathbb{Z}$ -linear combination of induced representations $\text{Ind}_{U_j}^G(\chi_j)$, as a virtual representation, where (U_j, V_j) is an element of $\mathfrak{F}$ and χ_j is a character of U_j/V_j of finite order.

In the following, we fix a family $\mathfrak{F}$ satisfying the hypothesis (b). For every $(U, V) \in \mathfrak{F}$, we have a homomorphism

$$\theta_{U,V}: K_1(\Lambda(G)) \longrightarrow \Lambda(U/V)^\times$$

which is the composite of the norm map of K -groups (see § 1.2)

$$\mathrm{Nr}_{\Lambda(G)/\Lambda(U)}: K_1(\Lambda(G)) \longrightarrow K_1(\Lambda(U))$$

and the canonical homomorphism

$$K_1(\Lambda(U)) \longrightarrow K_1(\Lambda(U/V)) = \Lambda(U/V)^\times.$$

Let S be the canonical Ore set for G . Then similarly we have a homomorphism

$$\theta_{S,U,V}: K_1(\Lambda(G)_S) \longrightarrow \Lambda(U/V)_S^\times.$$

Here we also denote the canonical Ore set for U/V by the same symbol S by abuse of notation.

Set

$$\theta = (\theta_{U,V})_{(U,V) \in \mathfrak{F}}: K_1(\Lambda(G)) \rightarrow \prod_{(U,V) \in \mathfrak{F}} \Lambda(U/V)^\times$$

and

$$\theta_S = (\theta_{S,U,V})_{(U,V) \in \mathfrak{F}}: K_1(\Lambda(G)_S) \rightarrow \prod_{(U,V) \in \mathfrak{F}} \Lambda(U/V)_S^\times.$$

Let Ψ_S be a subgroup of $\prod_{(U,V) \in \mathfrak{F}} \Lambda(U/V)_S^\times$ and let

$$\Psi = \Psi_S \cap \prod_{(U,V) \in \mathfrak{F}} \Lambda(U/V)^\times.$$

Definition 3.3 (The theta map, [Kato1] §2.4). Let G , $\mathfrak{F}$, θ_S , θ , Ψ_S and Ψ be as above.

If θ and θ_S satisfy

(θ -1) $\mathrm{Image}(\theta_S) \subseteq \Psi_S$,

(θ -2) $\mathrm{Image}(\theta) = \Psi$,

we call the induced surjective homomorphism

$$\theta: K_1(\Lambda(G)) \longrightarrow \Psi$$

the theta map for the group G , and call the induced homomorphism

$$\theta_S: K_1(\Lambda(G)_S) \longrightarrow \Psi_S$$

the localized theta map for the group G .

For $(U, V) \in \mathfrak{F}$, let F_U (resp. F_V) be the maximal subgroup of F^∞ fixed by U (resp. V). Since $\mathrm{Gal}(F_V/F_U) \cong U/V$ is abelian, the p -adic zeta function (pseudomeasure) $\xi_{U,V} \in \mathrm{Frac}(\Lambda(U/V))$ for F_V/F_U (see Remark 2.16) exists.

Theorem 3.4 (Burns, [Kato1] Proposition 2.5). *Let G be a compact p -adic Lie group. Assume that the theta map θ and its localized version θ_S for G exist, and also assume that $(\xi_{U,V})_{(U,V) \in \mathfrak{F}}$ is contained in Ψ_S .*

Then the p -adic zeta function $\xi_{F^\infty/F}$ for F^∞/F exists and satisfies the main conjecture.

Moreover, if θ is injective, $\xi_{F^\infty/F}$ is determined uniquely.

Proof. Let $f \in K_1(\Lambda(G)_S)$ be an arbitrary characteristic element for F^∞/F . Significantly, f is an element of $K_1(\Lambda(G)_S)$ which satisfies $\partial(f) = -[C_{F^\infty/F}]$.

Put $f_{U,V} = \theta_{S,U,V}(f)$ and $[C_{U,V}] = \theta_{U,V}([C])$ for each $(U,V) \in \mathfrak{F}$. Set $u_{U,V} = \xi_{U,V} f_{U,V}^{-1}$. Since the Iwasawa main conjecture is true for abelian extensions of totally real fields ([Wiles]), we have $\partial(\xi_{U,V}) = -[C_{U,V}]$. Hence we obtain $\partial(u_{U,V}) = 0$. By the localization exact sequence (Theorem 1.22), we have $u_{U,V} \in \Lambda(U/V)^\times$.

On the other hand, since $(f_{U,V})_{(U,V) \in \mathfrak{F}} \in \text{Image}(\theta_S) \subseteq \Psi_S$ by the condition $(\theta-1)$ and $(\xi_{U,V})_{(U,V) \in \mathfrak{F}} \in \Psi_S$ by assumption, $(u_{U,V})_{(U,V) \in \mathfrak{F}}$ is an element of Ψ_S . Therefore by the definition of Ψ , we have

$$(u_{U,V})_{(U,V) \in \mathfrak{F}} \in \Psi = \Psi_S \cap \prod_{(U,V) \in \mathfrak{F}} \Lambda(U/V)^\times.$$

By the condition $(\theta-2)$, there exists an element u of $K_1(\Lambda(G))$ which satisfies $\theta_{U,V}(u) = u_{U,V}$. We denote the image of u in $K_1(\Lambda(G)_S)$ by the same symbol u . Put $\xi_{F^\infty/F} = uf$. Since u is an element of $K_1(\Lambda(G))$, we have $\partial(u) = 0$. Therefore

$$\partial(\xi_{F^\infty/F}) = \partial(uf) = \partial(u) + \partial(f) = \partial(f) = -[C].$$

This implies that $\xi_{F^\infty/F}$ is also a characteristic element of $[C]$.

By the construction of $\xi_{F^\infty/F}$, it is clear that

$$(3.1) \quad \theta_{S,U,V}(\xi_{F^\infty/F}) = \xi_{U,V}.$$

Finally, we prove that $\xi_{F^\infty/F}$ satisfies the interpolation property (2.2). Since $\xi_{U,V}$ is the p -adic zeta function for F_V/F_U , it satisfies the following interpolating property:

$$(3.2) \quad \xi_{U,V}(\chi \otimes \kappa^r) = L_\Sigma(1-r; F_V/F_U, \chi) \quad \text{for } r \in \mathbb{N}, (p-1) \mid r$$

where χ is an arbitrary character of finite index of U/V .

By the hypothesis (b), an arbitrary Artin representation ρ is written in the form

$$\rho = \sum_{(U,V) \in \mathfrak{F}} \sum_{i \in I_{U,V}} a_{U,V}^{(i)} \text{Ind}_U^G(\chi_{U,V}^{(i)})$$

where $I_{U,V}$ is a index set, $a_{U,V}^{(i)}$ is an integer (assume all but finitely many $a_{U,V}^{(i)}$ are zero), and $\chi_{U,V}^{(i)}$ is a character of finite index of U/V . Then we have

$$\begin{aligned}
\xi_{F^\infty/F}(\rho \otimes \kappa^r) &= \prod_{(U,V) \in \mathfrak{F}} \prod_{i \in I_{U,V}} \xi(\text{Ind}_U^G(\chi_{U,V}^{(i)} \otimes \kappa^r)^{a_{U,V}^{(i)}}) \\
&= \prod_{(U,V) \in \mathfrak{F}} \prod_{i \in I_{U,V}} \theta_{S,U,V}(\xi)(\chi_{U,V}^{(i)} \otimes \kappa^r)^{a_{U,V}^{(i)}} \\
&= \prod_{(U,V) \in \mathfrak{F}} \prod_{i \in I_{U,V}} \xi_{U,V}(\chi_{U,V}^{(i)} \otimes \kappa^r)^{a_{U,V}^{(i)}} \quad (\text{by (3.1)}) \\
&= \prod_{(U,V) \in \mathfrak{F}} \prod_{i \in I_{U,V}} L_\Sigma(1-r; F_V/F_U, \chi_{U,V}^{(i)})^{a_{U,V}^{(i)}} \quad (\text{by (3.2)}) \\
&= L_\Sigma \left(1-r; F^\infty/F, \sum_{(U,V) \in \mathfrak{F}} \sum_i a_{U,V}^{(i)} \text{Ind}_U^G(\chi_{U,V}^{(i)}) \right) \\
&= L_\Sigma(1-r; F^\infty/F, \rho),
\end{aligned}$$

here the first equality follows from the definition of the evaluation maps, and the fifth equality follows from the compatibility of Artin L -functions with direct sum of representations.

The second equality follows from the next lemma (Lemma 3.5).

Note that if θ is injective, the element u above is determined uniquely, so is $\xi_{F^\infty/F}$. $\square$

Lemma 3.5. *Let (U, V) is an element of $\mathfrak{F}$. Then for an arbitrary character χ of U/V , the following diagram commutes.*

$$\begin{array}{ccc}
K_1(\Lambda(G)_S) & \xrightarrow{\theta_{S,U,V}} & K_1(\Lambda(U/V)_S) = \Lambda(U/V)_S^\times \\
\text{ev}_{\text{Ind}_U^G(\chi)} \downarrow & & \swarrow \text{ev}_\chi \\
\overline{\mathbb{Q}}_p \cup \{\infty\} & &
\end{array}$$

Proof. We will show the integral version of this lemma. In other words, we will prove the commutativity of the following diagram:

$$\begin{array}{ccc}
K_1(\Lambda(G)) & \xrightarrow{\theta_{U,V}} & K_1(\Lambda(U/V)) = \Lambda(U/V)^\times \\
\text{ev}_{\text{Ind}_U^G(\chi)} \downarrow & & \swarrow \text{ev}_\chi \\
\overline{\mathbb{Q}}_p^\times & &
\end{array}$$

Let W be the representation space of $\text{Ind}_U^G(\chi)$ over $\overline{\mathbb{Q}}$ and let W' be that of χ . Suppose that $\text{End}_{\overline{\mathbb{Q}}}(W)$ (resp. $\text{End}_{\overline{\mathbb{Q}}}(W')$) acts on W (resp. W') from the right.

Then by the definition of evaluation map, we obtain

$$\text{ev}_{\text{Ind}_U^G(\chi)} = [W \otimes_{\Lambda(G)} -].$$

On the other hand, we have

$$W \cong W' \otimes_{\Lambda(U/V)} \Lambda(G)$$

by the definition of induced representations, therefore we have

$$\mathrm{ev}_{\mathrm{Ind}_U^G(\chi)} = [W' \otimes_{\Lambda(U/V)} (\Lambda(U/V)\Lambda(G)_{\Lambda(G)}) \otimes_{\Lambda(G)} -],$$

and the right hand side is nothing but the definition of $\mathrm{ev}_\chi \circ \theta_{U,V}$.

It is not difficult to generalize this result to the localized version. $\square$

Kazuya Kato has constructed the theta maps for p -adic Lie groups of Heisenberg type ([Kato1], see also §8.1) and for certain open subgroups of $\mathbb{Z}_p^\times \ltimes \mathbb{Z}_p$ ([Kato2]). Kakde also constructed the theta map for the p -adic Lie group $H \rtimes \Gamma$ of “special type” where H is a compact pro- p abelian p -adic Lie group. He used the method of Kato in [Kato1] (See [Kakde]).

In our case $G = G^f \times \Gamma$, it is difficult to show that $(\xi_{U,V})_{(U,V) \in \mathfrak{F}} \in \Psi_S$ (See §7 and §8), so we should modify the proof of this proposition to show our main theorem (Theorem 3.1). But this technique gives us the ideas how to reduce the difficulties come from non-commutativity to the conditions of commutative cases.

4. THE ADDITIVE THETA MAP

In the following three sections, we construct the theta map (and its localized version) for the group $G = G^f \times \Gamma$. First, we construct a family $\mathfrak{F} = \{(U_i, V_i)\}$ which satisfies the hypothesis (b) of §3.2. Then we define a $\mathbb{Z}_p$ -module homomorphism

$$\theta^+ : \mathbb{Z}_p[[\mathrm{Conj}(G)]] \longrightarrow \prod_i \mathbb{Z}_p[[U_i/V_i]]$$

and characterize its image Ω . We show that θ^+ induces an isomorphism from $\mathbb{Z}_p[[\mathrm{Conj}(G)]]$ to Ω , which we call *the additive theta map for the group G* .

4.1. Construction of the family $\mathfrak{F}$. First, we define subgroups H and N of G^f as follows:

$$H = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & 0 \\ 0 & 1 & \mathbb{F}_p & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Note that N is abelian and normal in G . There are canonical isomorphisms

$$H \xrightarrow{\cong} \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix}; \begin{pmatrix} 1 & a & d & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & a & d \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix},$$

$$N \xrightarrow{\cong} \begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ \mathbb{F}_p \end{pmatrix}; \begin{pmatrix} 1 & 0 & 0 & f \\ 0 & 1 & 0 & e \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} f \\ e \\ c \end{pmatrix}.$$

By the isomorphisms above, we identify H with $\begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix}$ and N with $\begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ \mathbb{F}_p \end{pmatrix}$.

Then G^f is represented as a semi-direct product

$$G^f \cong H \ltimes N = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix} \ltimes \begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ \mathbb{F}_p \end{pmatrix},$$

where H acts on N from the left as ordinary product of matrices.

From representation theory of semi-direct products of finite groups (See [Serre1], Chapitre 8.2), all irreducible representations of G^f are obtained from representations of H and those of N . Let us review this construction.

First, let $\mathfrak{X}(N)$ be the character group of the abelian group N :

$$\mathfrak{X}(N) = \left\{ \chi_{ijk} \mid 0 \leq i, j, k \leq p-1 \right\}$$

where $\chi_{ijk} \left(\begin{pmatrix} f \\ e \\ c \end{pmatrix} \right) = \exp \left(\frac{2\pi\sqrt{-1}}{p} (fi + ej + ck) \right) \quad (c, e, f \in \mathbb{F}_p)$.

The group H acts on $\mathfrak{X}(N)$ from the right by

$$(\chi * h)(n) = \chi(hn) \quad \text{for all } n \in N$$

where $h \in H$ and $\chi \in \mathfrak{X}(N)$. Specifically,

$$\chi_{ijk} * \begin{pmatrix} 1 & a & d \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} = \chi_{i, ai+j, di+bj+k} \quad \text{for every} \quad \begin{pmatrix} 1 & a & d \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \in H$$

where $\chi_{ijk} \in \mathfrak{X}(N)$. Then we have

$$\chi_{00k} \quad (0 \leq k \leq p-1), \quad \chi_{0j0} \quad (1 \leq j \leq p-1), \quad \chi_{i00} \quad (1 \leq i \leq p-1)$$

as a system of representatives for the orbital decomposition $\mathfrak{X}(N)/H$.

Next, let H_{ijk} be the isotropic subgroup of H at each χ_{ijk} . Then for each representative above, we have

$$\begin{aligned} H_{00k} &= H_0 = H \quad (0 \leq k \leq p-1), \\ H_{0j0} &= H_1 = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (1 \leq j \leq p-1), \\ H_{i00} &= H_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix} \quad (1 \leq i \leq p-1). \end{aligned}$$

Let $G_\ell^f = H_\ell \ltimes N$ for $\ell = 0, 1, 2$. Then we may extend the domain of the character χ_{00k} (resp. χ_{0j0} , χ_{i00}) to G_0^f (resp. G_1^f , G_2^f) by setting

$$\chi_{00k}(h_0 n) = \chi_{00k}(n), \quad \chi_{0j0}(h_1 n) = \chi_{0j0}(n), \quad \chi_{i00}(h_2 n) = \chi_{i00}(n)$$

where $h_\ell \in H_\ell$ ($\ell = 0, 1, 2$) and $n \in N$. Note that each χ_{00k} (resp. χ_{0j0} , χ_{i00}) is a character of degree 1 of the group G_0^f (resp. G_1^f , G_2^f).

Now let ρ_ℓ ($\ell = 0, 1, 2$) be an arbitrary irreducible representation of H_ℓ . Then we obtain an irreducible representation of G_ℓ^f by composing ρ_ℓ with the canonical projection $G_\ell^f \rightarrow H_\ell$, which we also denote by ρ_ℓ . Consider the tensor products of representations $\chi_{00k} \otimes \rho_0$ (resp. $\chi_{0j0} \otimes \rho_1$, $\chi_{i00} \otimes \rho_2$), and let

$$\begin{aligned}\theta_{k,\rho_0} &= \text{Ind}_{G_0^f}^{G^f}(\chi_{00k} \otimes \rho_0) = \chi_{00k} \otimes \rho_0, \\ \theta_{j,\rho_1} &= \text{Ind}_{G_1^f}^{G^f}(\chi_{0j0} \otimes \rho_1), \\ \theta_{i,\rho_2} &= \text{Ind}_{G_2^f}^{G^f}(\chi_{i00} \otimes \rho_2).\end{aligned}$$

Proposition 4.1. *Suppose $0 \leq k \leq p-1$, $1 \leq j \leq p-1$ and $1 \leq i \leq p-1$. Let ρ_ℓ be an irreducible representation of H_ℓ . Then each θ_{k,ρ_0} , θ_{j,ρ_1} , θ_{i,ρ_2} is an irreducible representation of G^f . Moreover, an arbitrary irreducible representation of G^f is isomorphic to one of the θ_{k,ρ_0} , θ_{j,ρ_1} , θ_{i,ρ_2} .*

Proof. See [Serre1], Chapitre 8.2 (for irreducibility of each θ , we use Mackey's irreducibility criterion). $\square$

Note that H_1 and H_2 are abelian groups, so if we set

$$\begin{aligned}U_1^f &= H_1 \ltimes N & V_1' &= \{I_3\} \ltimes \begin{pmatrix} \mathbb{F}_p \\ 0 \\ \mathbb{F}_p \end{pmatrix} \\ &= \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, & &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ U_2^f &= H_2 \ltimes N & V_2' &= \{I_3\} \ltimes \begin{pmatrix} 0 \\ \mathbb{F}_p \\ \mathbb{F}_p \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, & &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix},\end{aligned}$$

each irreducible representation θ_{j,ρ_1} (resp. θ_{i,ρ_2}) is obtained as the induced representation $\text{Ind}_{U_1^f}^{G^f}(\chi_1')$ (resp. $\text{Ind}_{U_2^f}^{G^f}(\chi_2')$) where χ_1' (resp. χ_2') is a certain character of U_1^f/V_1' (resp. U_2^f/V_2') of finite order.

On the other hand, we may regard $H_0(=H)$ as a semi-direct product

$$H \cong \begin{pmatrix} 1 & \mathbb{F}_p \\ 0 & 1 \end{pmatrix} \ltimes (\mathbb{F}_p \quad \mathbb{F}_p); \begin{pmatrix} 1 & a & d \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \ltimes (a \quad d)$$

where $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in \begin{pmatrix} 1 & \mathbb{F}_p \\ 0 & 1 \end{pmatrix}$ acts on $(a \ d) \in (\mathbb{F}_p \ \mathbb{F}_p)$ from the left by

$$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} * (a \ d) = (a \ d) \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}^{-1} \quad (\text{ordinary product of matrices}).$$

Let

$$\begin{aligned} U_0^H &= H & V_0^H &= \{I_2\} \ltimes (0 \ \mathbb{F}_p) \\ &= \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix}, & &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ U_1^H &= \{I_2\} \ltimes (\mathbb{F}_p \ \mathbb{F}_p) & V_1^H &= \{I_2\} \ltimes (0 \ 0) \\ &= \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & &= \{I_3\}, \end{aligned}$$

then by an argument similar to above, every irreducible representation of H is obtained as the induced representation $\text{Ind}_{U_\ell^H}^H(\chi_\ell)$ ($\ell = 0$ or 1) where χ_ℓ is a certain character of U_ℓ^H/V_ℓ^H . Therefore if we set

$$\begin{aligned} U_0^f &= U_0^H \ltimes N & V_0^f &= V_0^H \ltimes \begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ 0 \end{pmatrix} \\ &= G, & &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ U_1^f &= U_1^H \ltimes N & V_1'' &= V_1^H \ltimes \begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, & &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \end{aligned}$$

each irreducible representation θ_{k,ρ_0} is obtained as the induced representation $\text{Ind}_{U_0^f}^{G^f}(\chi_0)$ (resp. $\text{Ind}_{U_1^f}^{G^f}(\chi_1'')$) where χ_0 (resp. χ_1'') is a certain character of U_0^f/V_0^f (resp. U_1^f/V_1'') of finite order.

Let $V_1^f = V_1' \cap V_1'' = \langle \zeta \rangle$. Then by the argument above, every irreducible representation of G^f is obtained as the induced representation of a certain character $\chi_i \in \mathfrak{X}(U_i^f/V_i^f)$ where $\mathfrak{X}(U_i^f/V_i^f)$ is the character group of the abelian group U_i^f/V_i^f . Hence $\mathfrak{F}^f = \{(U_0^f, V_0^f), (U_1^f, V_1^f), (U_2^f, V_2^f)\}$ satisfies the hypothesis (b) for the group G^f .

For certain technical reasons, we replace V_2' by

$$V_2^f = \{I_3\} \ltimes \begin{pmatrix} 0 \\ \mathbb{F}_p \\ \mathbf{0} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbf{0} \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

and add following subgroups to our family $\mathfrak{F}^f$

$$\begin{aligned} \widetilde{U}_2^f &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix} \ltimes N & \quad \widetilde{V}_2^f &= \{I_3\} \ltimes \begin{pmatrix} \mathbb{F}_p \\ \mathbb{F}_p \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \quad &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ U_3^f &= \{I_3\} \ltimes N & \quad V_3^f &= \{I_3\} \ltimes \{0\} \\ &= \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \quad &= \{I_4\}. \end{aligned}$$

Note that V_2^f (resp. $\widetilde{V}_2^f, V_3^f$) is normal in U_2^f (resp. $\widetilde{U}_2^f, U_3^f$) and U_2^f/V_2^f (resp. $\widetilde{U}_2^f/\widetilde{V}_2^f, U_3^f/V_3^f$) is abelian.

For each i (including $\widetilde{U}_2$ and $\widetilde{V}_2$), let

$$U_i = U_i^f \times \Gamma, \quad V_i = V_i^f \times \{1\}$$

and let $\mathfrak{F} = \{(U_i, V_i)\}_i$. It is clear that $\mathfrak{F}$ satisfies the hypothesis (b) for the group G .

4.2. Construction of the isomorphism θ^+ . In the following, we denote

$$\begin{aligned} \prod_i \mathbb{Z}_p[[U_i/V_i]] &= \mathbb{Z}_p[[U_0/V_0]] \times \mathbb{Z}_p[[U_1/V_1]] \\ &\quad \times \mathbb{Z}_p[[\widetilde{U}_2/\widetilde{V}_2]] \times \mathbb{Z}_p[[U_2/V_2]] \times \mathbb{Z}_p[[U_3/V_3]] \end{aligned}$$

and we use the notation U_i (resp. V_i) for one of the subgroups $U_0, U_1, \widetilde{U}_2, U_2$ and U_3 (resp. $V_0, V_1, \widetilde{V}_2, V_2$ and V_3).

For an arbitrary finite group F , we denote by $\mathbb{Z}_p[\text{Conj}(F)]$ the free $\mathbb{Z}_p$ -module of finite rank with free bases $\text{Conj}(F)$. For an arbitrary pro-finite group P , let $\mathbb{Z}_p[[\text{Conj}(P)]]$ be the projective limit of the free $\mathbb{Z}_p$ -modules $\mathbb{Z}_p[\text{Conj}(P_\lambda)]$ over finite quotient groups P_λ of P .

Let $\{u_1, u_2, \dots, u_{r_i}\} \subseteq G$ be a system of representatives of the coset decomposition G/U_i for each i . Then for an arbitrary conjugacy class $[g] \in \text{Conj}(G)$, set

$$\text{Tr}_i([g]) = \sum_{j=1}^{r_i} \tau_j([g])$$

where $\tau_j([g]) = [u_j^{-1}gu_j]$ if $u_j^{-1}gu_j$ is contained in U_i , and $\tau_j([g]) = 0$ otherwise. It is easy to see that $\text{Tr}_i([g])$ is independent of the choice of the

representatives $\{u_j\}_{j=1}^{r_i}$, therefore Tr_i induces a well-defined $\mathbb{Z}_p$ -module homomorphism

$$\text{Tr}_i: \mathbb{Z}_p[[\text{Conj}(G)]] \longrightarrow \mathbb{Z}_p[[\text{Conj}(U_i)]],$$

which we call *the trace homomorphism from $\mathbb{Z}_p[[\text{Conj}(G)]]$ to $\mathbb{Z}_p[[\text{Conj}(U_i)]]$* .

We define the homomorphism θ_i^+ as the composition of the trace map Tr_i and the natural surjection

$$\mathbb{Z}_p[[\text{Conj}(U_i)]] \longrightarrow \mathbb{Z}_p[[U_i/V_i]].$$

Now let us calculate $\theta_i^+([g_0])$ for $[g_0] \in \text{Conj}(G^f)$. Here we regard $[g_0]$ as an element $([g_0], 0)$ contained in $\text{Conj}(G) = \text{Conj}(G^f) \times \text{Conj}(\Gamma)$.

Here we only compute $\theta_2^+([g_0])$. We may take $\{\alpha^j \delta^k \mid 0 \leq j, k \leq p-1\}$ as a system of representatives of G/U_2 . If $g_0 = \begin{pmatrix} 1 & a & d & f \\ 0 & 1 & b & e \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}$, we have

$$(\alpha^j \delta^k)^{-1} g_0 (\alpha^j \delta^k) = \begin{pmatrix} 1 & a & d - bj & f - ej - ck \\ 0 & 1 & b & e \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

This implies that $(\alpha^j \delta^k)^{-1} g_0 (\alpha^j \delta^k)$ is contained in U_2 if and only if $a = 0$ and $d - bj = 0$.

First, suppose that $a = 0$ and $b \neq 0$. Then there exists a unique j satisfying $d - bj = 0$. We denote this j by $\frac{d}{b}$. Then we have

$$\begin{aligned} \theta_2^+([g_0]) &= \sum_{k=0}^{p-1} \begin{pmatrix} 1 & 0 & 0 & f - e \cdot \frac{d}{b} - ck \\ 0 & 1 & b & e \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix} \pmod{\langle \varepsilon \rangle} \\ &= \begin{cases} \beta^b \gamma^c (1 + \zeta + \dots + \zeta^{p-1}) & \text{if } c \neq 0, \\ p \beta^b \zeta^{f - e \cdot \frac{d}{b}} & \text{if } c = 0. \end{cases} \end{aligned}$$

If $a = b = 0$ and $d \neq 0$, $(\alpha^j \delta^k)^{-1} g_0 (\alpha^j \delta^k)$ is not contained in U_2 for each $0 \leq j, k \leq p-1$, hence $\theta_2^+([g_0]) = 0$.

If $a = b = d = 0$, we have

$$\begin{aligned} \theta_2^+([g_0]) &= \sum_{j,k=0}^{p-1} \begin{pmatrix} 1 & 0 & 0 & f - ej - ck \\ 0 & 1 & 0 & e \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix} \pmod{\langle \varepsilon \rangle} \\ &= \begin{cases} p^2 \zeta^f & \text{if } c = e = 0, \\ p \gamma^c (1 + \zeta + \dots + \zeta^{p-1}) & \text{if } c \neq 0, \\ p(1 + \zeta + \dots + \zeta^{p-1}) & \text{if } c = 0, e \neq 0. \end{cases} \end{aligned}$$

In this way, we have

$$\theta_2^+([g_0]) = \begin{cases} 0 & \text{if } a \neq 0 \text{ or } a = b = 0, d \neq 0, \\ \beta^b \gamma^c (1 + \zeta + \dots + \zeta^{p-1}) & \text{if } a = 0, b \neq 0, c \neq 0, \\ p\beta^b \zeta^{f - e \cdot \frac{d}{b}} & \text{if } a = c = 0, b \neq 0, \\ p\gamma^c (1 + \zeta + \dots + \zeta^{p-1}) & \text{if } a = b = d = 0, c \neq 0, \\ p(1 + \zeta + \dots + \zeta^{p-1}) & \text{if } a = b = c = d = 0, e \neq 0, \\ p^2 \zeta^f & \text{if } a = b = c = d = e = 0. \end{cases}$$

Similarly, we may take $\{\beta^j \mid 0 \leq j \leq p-1\}$ for a system of representatives of G/U_1 , $\{\alpha^j \mid 0 \leq j \leq p-1\}$ for that of $G/\widetilde{U}_2$, and $\{\alpha^j \beta^k \delta^\ell \mid 0 \leq j, k, \ell \leq p-1\}$ for that of G/U_3 . Using these representatives, we may calculate $\theta^+([g_0])$ as follows:

$$\theta^+ : \mathbb{Z}_p[[\text{Conj}(G)]] \longrightarrow \prod_i \mathbb{Z}_p[[U_i/V_i]]$$

$$\mathbf{c}_I(a, b, c) = \{\alpha^a \beta^b \gamma^c \delta^d \varepsilon^e \zeta^f \mid 0 \leq d, e, f \leq p-1\} \quad (a \neq 0, b \neq 0)$$

$$\mapsto (\alpha^a \beta^b \gamma^c, 0, 0, 0, 0),$$

$$\mathbf{c}_{II}(a, c, d, e) = \{\alpha^a \gamma^c \delta^d \varepsilon^e \zeta^f \mid 0 \leq f \leq p-1\} \quad (a \neq 0, c \neq 0)$$

$$\mapsto (\alpha^a \gamma^c, \alpha^a \gamma^c \delta^d \varepsilon^e h_{\varepsilon^c \delta^{-a}}, 0, 0, 0),$$

$$\mathbf{c}_{III}(a, e) = \{\alpha^a \varepsilon^e \delta^d \zeta^f \mid 0 \leq d, f \leq p-1\} \quad (a \neq 0)$$

$$\mapsto (\alpha^a, \alpha^a \varepsilon^e h_\delta, 0, 0, 0),$$

$$\mathbf{c}_{IV}(b, c) = \{\beta^b \gamma^c \delta^d \varepsilon^e \zeta^f \mid 0 \leq d, e, f \leq p-1\} \quad (b \neq 0, c \neq 0)$$

$$\mapsto (\beta^b \gamma^c, 0, \beta^b \gamma^c h_\delta, \beta^b \gamma^c h_\zeta, 0),$$

$$\mathbf{c}_V(b, f) = \{\beta^b \delta^d \varepsilon^e \zeta^{f'} \mid f' - \frac{d}{b}e \equiv f \pmod{p}\} \quad (b \neq 0)$$

$$\mapsto (\beta^b, 0, \beta^b h_\delta, p\beta^b \zeta^f, 0),$$

$$\mathbf{c}_{VI}(c, d) = \{\gamma^c \delta^d \varepsilon^e \zeta^f \mid 0 \leq e, f \leq p-1\} \quad (c \neq 0, d \neq 0)$$

$$\mapsto (\gamma^c, \gamma^c \delta^d h_\varepsilon, p\gamma^c \delta^d, 0, 0),$$

$$\mathbf{c}_{VII}(c) = \{\gamma^c \varepsilon^e \zeta^f \mid 0 \leq e, f \leq p-1\} \quad (c \neq 0)$$

$$\mapsto (\gamma^c, \gamma^c h_\varepsilon, p\gamma^c, p\gamma^c h_\zeta, p\gamma^c h_\varepsilon h_\zeta),$$

$$\mathbf{c}_{VIII}(d, e) = \{\delta^d \varepsilon^e \zeta^f \mid 0 \leq f \leq p-1\} \quad (d \neq 0)$$

$$\mapsto (1, p\delta^d \varepsilon^e, p\delta^d, 0, 0)$$

$$\mathbf{c}_{IX}(e) = \{\varepsilon^e \zeta^f \mid 0 \leq f \leq p-1\} \quad (e \neq 0)$$

$$\mapsto (1, p\varepsilon^e, p, ph_\zeta, p^2 \varepsilon^e h_\zeta),$$

$$\mathbf{c}_X(f) = \{\zeta^f\}$$

$$\mapsto (1, p, p, p^2 \zeta^f, p^3 \zeta^f),$$

where for an element $g_0 \in G^f$, h_{g_0} denotes an element $1 + g_0 + g_0^2 + \dots + g_0^{p-1}$ of $\mathbb{Z}_p[[\text{Conj}(G)]]$.

Now let us consider the finite quotients $G^{(n)} = G^f \times \Gamma/\Gamma^{p^n}$ and $U_i^{(n)} = U_n^f \times \Gamma/\Gamma^{p^n}$. Then we have

$$\begin{aligned}\mathbb{Z}_p[G^{(n)}] &\cong \mathbb{Z}_p[G^f] \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\Gamma/\Gamma^{p^n}], \\ \mathbb{Z}_p[U_i^{(n)}/V_i] &\cong \mathbb{Z}_p[U_i^f/V_i^f] \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\Gamma/\Gamma^{p^n}].\end{aligned}$$

θ_i^+ induces a homomorphism

$$\theta_i^{+, (n)}: \mathbb{Z}_p[G^{(n)}] \rightarrow \mathbb{Z}_p[U_i^{(n)}/V_i]$$

which sends $[g] = ([g_0], t^z) \in \text{Conj}(G^{(n)}) = \text{Conj}(G^f) \times \Gamma/\Gamma^{p^n}$ to $(\theta_i^+([g_0]), t^z)$. For example, we have

$$\theta^{+, (n)}((c_I(a, b, c), t^z)) = ((\alpha^a \beta^b \gamma^c, t^z), (0, t^z), (0, t^z), (0, t^z), (0, t^z)),$$

and so on.

We would like to characterize the image of $\theta^{+, (n)}$.

Let I_i^f be the submodule of $\mathbb{Z}_p[U_i^f/V_i^f]$ defined as follows:

$$\begin{aligned}I_1^f &= [p\delta^d \varepsilon^e, \alpha^a \varepsilon^e h_\delta (a \neq 0), \gamma^c \delta^d h_\varepsilon (c \neq 0), \alpha^a \gamma^c \delta^d \varepsilon^e h_{\varepsilon^c \delta^{-a}} (a \neq 0, c \neq 0)]_{\mathbb{Z}_p}, \\ \tilde{I}_2^f &= [\beta^b \gamma^c h_\delta (b \neq 0), p\gamma^c \delta^d]_{\mathbb{Z}_p}, \\ I_2^f &= [p^2 \zeta^f, p\gamma^c h_\zeta, \beta^b \gamma^c h_\zeta (b \neq 0, c \neq 0), p\beta^b \zeta^f (b \neq 0)]_{\mathbb{Z}_p}, \\ I_3^f &= [p^3 \zeta^f, p^2 \varepsilon^e h_\zeta (e \neq 0), p\gamma^c h_\varepsilon h_\zeta (c \neq 0)]_{\mathbb{Z}_p}.\end{aligned}$$

Here we denote by $[S]_R$ the R -submodule of M generated by S over R where R is a commutative ring, M is an R -module and S is a finite subset of M . Each I_i^f is a $\mathbb{Z}_p$ -submodule of $\mathbb{Z}_p[U_i^f/V_i^f]$ generated by $\{\theta_i^+([g_0]) \mid g_0 \in G^f\}$.

Let

$$I_i^{(n)} = I_i^f \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\Gamma/\Gamma^{p^n}] \quad \left(\subseteq \mathbb{Z}_p[U_i^{(n)}/V_i] \right)$$

and

$$I_i = \varprojlim_n I_i^{(n)} \quad (\subseteq \mathbb{Z}_p[[U_i/V_i]]).$$

For each i , I_i is the image of the homomorphism θ_i^+ .

Note that generators of $I_1^{(n)}$ (resp. $\tilde{I}_2^{(n)}$, $I_3^{(n)}$) above are $\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]$ -linearly independent, but those of $I_2^{(n)}$ are not $\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]$ -linearly independent, which have a relation

$$(4.1) \quad \sum_{f=0}^{p-1} p^2 \zeta^f = p \cdot p h_\zeta.$$

Definition 4.2. We define Ω to be the $\mathbb{Z}_p$ -submodule of $\prod_i \mathbb{Z}_p[[U_i/V_i]]$ consisting of all elements $(y_0, y_1, \tilde{y}_2, y_2, y_3)$ satisfying the following two conditions:

- (1) (trace relations)
 - (rel-1) $\text{Tr}_{\mathbb{Z}_p[[U_0/V_0]]/\mathbb{Z}_p[[U_1/V_0]]} y_0 \equiv y_1,$
 - (rel-2) $\text{Tr}_{\mathbb{Z}_p[[U_0/V_0]]/\mathbb{Z}_p[[\tilde{U}_2/V_0]]} y_0 \equiv \tilde{y}_2,$
 - (rel-3) $\text{Tr}_{\mathbb{Z}_p[[\tilde{U}_2/\tilde{V}_2]]/\mathbb{Z}_p[[U_2/\tilde{V}_2]]} \tilde{y}_2 \equiv y_2,$




FIGURE 1. Trace and norm relations.

$$(\text{rel-4}) \quad \text{Tr}_{\mathbb{Z}_p[[U_1/\widetilde{V}_2]]/\mathbb{Z}_p[[U_1 \cap \widetilde{U}_2/\widetilde{V}_2]]} y_1 \equiv \text{Tr}_{\mathbb{Z}_p[[\widetilde{U}_2/\widetilde{V}_2]]/\mathbb{Z}_p[[U_1 \cap \widetilde{U}_2/\widetilde{V}_2]]} \widetilde{y}_2,$$

$$(\text{rel-5}) \quad \text{Tr}_{\mathbb{Z}_p[[U_1/V_1]]/\mathbb{Z}_p[[U_3/V_1]]} y_1 \equiv y_3,$$

$$(\text{rel-6}) \quad \text{Tr}_{\mathbb{Z}_p[[U_2/V_2]]/\mathbb{Z}_p[[U_3/V_2]]} y_2 \equiv y_3.$$

(These trace relations are described in Figure 1.)

(2) $y_i \in I_i$ for each i .

Proposition-Definition 4.3. *The homomorphism $\theta^+ = (\theta_i^+)$ induces an isomorphism*

$$\theta^+ : \mathbb{Z}_p[[\text{Conj}(G)]] \xrightarrow{\cong} \Omega.$$

We call this induced isomorphism θ^+ *the additive theta map for G* .

Proof. It is clear from the calculation above that Ω contains the image of θ^+ . Hence it is sufficient to prove its injectivity and surjectivity.

We now show that θ^+ induces an isomorphism on the finite quotients

$$\theta^{+, (n)} : \mathbb{Z}_p[\text{Conj}(G^{(n)})] \xrightarrow{\cong} \Omega^{(n)},$$

for every $n \geq 1$, where $\Omega^{(n)}$ is defined to be the subgroup of $\prod_i \mathbb{Z}_p[U_i^{(n)}/V_i]$

by the same conditions in Definition 4.2. Then by taking the projective limit, we obtain the desired isomorphism.

To simplify the notation, we denote the induced homomorphism $\theta^{+, (n)}$ by θ^+ in the following.

- Injectivity.

Let $y \in \mathbb{Z}_p[\text{Conj}(G^{(n)})]$ be an element satisfying $\theta^+(y) = 0$. Then from the hypothesis (b) for the family $\{(U_i^{(n)}, V_i)\}_i$,⁵ an arbitrary

⁵Recall that an arbitrary irreducible representation τ of $G^{(n)} = G^f \times \Gamma/\Gamma^{p^n}$ is isomorphic to $\rho \otimes \omega_n$ where ρ is an irreducible representation of G^f and ω_n is a character of Γ/Γ^{p^n} . By the construction of U_i and V_i , an arbitrary irreducible representation ρ of G^f is described as $\sum_{i,j} a_i^{(j)} \text{Ind}_{U_i^f}^{G^f}(\chi_i^{(j)})$ where $a_i^{(j)}$ is an integer and $\chi_i^{(j)}$ is a character

irreducible representation of $G^{(n)}$ is isomorphic to a $\mathbb{Z}$ -linear combination (as a virtual representation) of $\text{Ind}_{U_i^{(n)}}^{G^{(n)}}(\chi_i)$'s where χ_i is a character of $U_i^{(n)}/V_i$. We denote by $\tilde{\chi}_i$ the character of the induced representation $\text{Ind}_{U_i^{(n)}}^{G^{(n)}}(\chi_i)$. Then we have

$$\tilde{\chi}_i(y) = \sum_{[g] \in \text{Conj}(G^{(n)})} m_{[g]} \sum_{j, u_j^{-1}gu_j \in U_i^{(n)}} \chi_i(u_j^{-1}gu_j) = \chi_i \circ \theta_i^+(y)$$

by the property of induced characters, where $\{u_1, \dots, u_{r_i}\}$ is a system of representatives of $G^{(n)}/U_i^{(n)}$ and

$$y = \sum_{[g] \in \text{Conj}(G^{(n)})} m_{[g]}[g] \quad (m_g \in \mathbb{Z}_p).$$

On the other hand, $\theta_i^+(y)$ vanishes by assumption, hence $\tilde{\chi}_i(y) = 0$. This implies that $\chi(y) = 0$ for an arbitrary irreducible character χ of $G^{(n)}$. Since all irreducible characters of $G^{(n)}$ form a dual basis of the $\overline{\mathbb{Q}}_p$ -vector space $\overline{\mathbb{Q}}_p \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[\text{Conj}(G^{(n)})]$, we may conclude that $y = 0$.

- Surjectivity.

Let $(y_0, y_1, \tilde{y}_2, y_2, y_3)$ be an arbitrary element of $\Omega^{(n)}$, then each y_i is a $\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]$ -linear combination of generators of $I_i^{(n)}$:

$$\begin{aligned} y_0 &= \sum_{a,b,c} \kappa_{abc} \alpha^a \beta^b \gamma^c, \\ y_1 &= \sum_{d,e} \lambda_{de}^{(1)} \delta^d \varepsilon^e + \sum_{a \neq 0,e} \lambda_{ae}^{(2)} \alpha^a \varepsilon^e h_\delta \\ &\quad + \sum_{c \neq 0,d} \lambda_{cd}^{(3)} \gamma^c \delta^d h_\varepsilon + \sum_{a \neq 0, c \neq 0, d, e} \lambda_{acde}^{(4)} \alpha^a \gamma^c \delta^d \varepsilon^e h_{\varepsilon^c \delta^d - a}, \\ \tilde{y}_2 &= \sum_{b \neq 0, c} \mu_{bc}^{(1)} \beta^b \gamma^c h_\delta + \sum_{c,d} p \mu_{cd}^{(2)} \gamma^c \delta^d, \\ y_2 &= \sum_f p^2 \nu_f^{(1)} \zeta^f + \sum_c p \nu_c^{(2)} \gamma^c h_\zeta \\ &\quad + \sum_{b \neq 0, c \neq 0} \nu_{bc}^{(3)} \beta^b \gamma^c h_\zeta + \sum_{b \neq 0, f} p \nu_{bf}^{(4)} \beta^b \zeta^f, \\ y_3 &= \sum_f p^3 \sigma_f^{(1)} \zeta^f + \sum_{e \neq 0} p^2 \sigma_e^{(2)} \varepsilon^e h_\zeta + \sum_{c \neq 0} p \sigma_c^{(3)} \gamma^c h_\varepsilon h_\zeta. \end{aligned}$$

Note that by the relation (4.1), $\nu_f^{(1)} (0 \leq f \leq p-1)$ and $\nu_0^{(2)}$ are not determined uniquely.

of U_i^f/V_i^f of finite index. Then we have $\tau = \rho \otimes \omega_n = \left(\sum_{i,j} a_i^{(j)} \text{Ind}_{U_i^f}^{G^f}(\chi_i^{(j)}) \right) \otimes \omega_n = \sum_{i,j} a_i^{(j)} \text{Ind}_{U_i^{(n)}}^{G^{(n)}}(\chi_i^{(j)} \otimes \omega_n)$. $\chi_{i,j} \otimes \omega_n$ is a character of $U_i^{(n)}/V_i$ of finite index, hence $\{U_i^{(n)}, V_i\}_i$ also satisfies the hypothesis (b) for the group $G^{(n)}$.

The trace relations in Definition 4.2 (1) give linear relations among these coefficients. For example, let us describe the (rel-6) explicitly. Take $\{\beta^j \mid 0 \leq j \leq p-1\}$ as a system of representatives of $U_2^{(n)}/U_3^{(n)}$. Then for $u_2 \in U_2^f/V_2^f$, $\beta^{-j}u_2\beta^j$ is contained in $U_3^{(n)}/V_2$ if and only if $b(u_2) = 0$ where

$$u_2 \equiv \beta^{b(u_2)}\gamma^{c(u_2)}\zeta^{f(u_2)} \pmod{\langle \varepsilon \rangle}.$$

It is obvious that $\beta^{-j}u_2\beta^j = u_2$ when $b(u_2) = 0$. Therefore,

$$\begin{aligned} \text{Tr}_{\mathbb{Z}_p[U_2^{(n)}/V_2]/\mathbb{Z}_p[U_3^{(n)}/V_2]} y_2 &= \sum_f p^2 (p\nu_f^{(1)} + \nu_0^{(2)})\zeta^f \\ &\quad + \sum_{c \neq 0} p^2 \nu_c^{(2)} \gamma^c h_\zeta. \end{aligned}$$

On the other hand,

$$y_3 \equiv p^2 \sum_{e \neq 0, f} (p\sigma_f^{(1)} + \sigma_e^{(2)})\zeta^f + \sum_{c \neq 0} p^2 \sigma_c^{(3)} \gamma^c h_\zeta \pmod{V_2}.$$

By comparing these coefficients, we have

$$(4.2) \quad p\nu_f^{(1)} + \nu_0^{(2)} = p\sigma_f^{(1)} + \sum_{e \neq 0} \sigma_e^{(2)} \quad (0 \leq f \leq p-1),$$

$$(4.3) \quad \nu_c^{(2)} = \sigma_c^{(3)} \quad (1 \leq c \leq p-1).$$

However, because of the relation (4.1), we may change $\nu_f^{(1)} \rightarrow \nu_f^{(1)} + z$ and $\nu_0^{(2)} \rightarrow \nu_0^{(2)} - pz$ for every $z \in \mathbb{Z}_p[\Gamma/\Gamma^{p^n}]$. Therefore in the equation (4.2), we may choose $\nu_f^{(1)}$ and $\nu_0^{(2)}$ as they satisfy

$$\begin{aligned} \nu_f^{(1)} &= \sigma_f^{(1)} \quad (0 \leq f \leq p-1), \\ \nu_0^{(2)} &= \sum_{e \neq 0} \sigma_e^{(2)}. \end{aligned}$$

Similarly, we can describe the other trace relations explicitly as follows:

$$(rel-1) \quad \kappa_{000} = \sum_{d,e} \lambda_{de}^{(1)}, \quad \kappa_{a00} = \sum_e \lambda_{ae}^{(2)} \quad (a \neq 0),$$

$$\kappa_{00c} = \sum_d \lambda_{cd}^{(3)} \quad (c \neq 0),$$

$$\kappa_{a0c} = \sum_{d,e} \lambda_{acde}^{(4)} \quad (a \neq 0, c \neq 0),$$

$$(rel-2) \quad \kappa_{00c} = \sum_d \mu_{cd}^{(2)}, \quad \kappa_{0bc} = \mu_{bc}^{(1)} \quad (b \neq 0),$$

$$(rel-3) \quad \mu_{bc}^{(1)} = \nu_{bc}^{(3)} \quad (b \neq 0, c \neq 0),$$

$$\mu_{b0}^{(1)} = \sum_f \nu_{bf}^{(4)} \quad (b \neq 0),$$

$$\mu_{c0}^{(2)} = \nu_c^{(2)} \quad (c \neq 0), \quad \mu_{00}^{(2)} = \sum_f \nu_f^{(1)} + \nu_0^{(2)},$$

$$\text{(rel-4)} \quad \sum \lambda_{de}^{(1)} = \mu_{0d}^{(2)}, \lambda_{cd}^{(3)} = \mu_{cd}^{(2)} \quad (c \neq 0)$$

$$\text{(rel-5)} \quad \lambda_{00}^{(1)} = \sum_f \sigma_f^{(1)}, \lambda_{0e}^{(1)} = \sigma_e^{(2)} \quad (e \neq 0),$$

$$\text{(rel-6)} \quad \lambda_{c0}^{(3)} = \sigma_c^{(3)} \quad (c \neq 0), \\ \nu_f^{(1)} = \sigma_f^{(1)}, \nu_0^{(2)} = \sum_{e \neq 0} \sigma_e^{(2)}, \nu_c^{(2)} = \sigma_c^{(3)} \quad (c \neq 0).$$

Then we may show by direct calculation, using the explicit trace relations above, that an element of $\mathbb{Z}_p[\text{Conj}(G^{(n)})]$

$$\begin{aligned} y = & \sum_{a \neq 0, b \neq 0, c} \kappa_{abc} \mathbf{c}_I(a, b, c) + \sum_{a \neq 0, c \neq 0, d, e} \lambda_{acde}^{(4)} \mathbf{c}_{II}(a, c, d, e) \\ & + \sum_{a \neq 0, e} \lambda_{ae}^{(2)} \mathbf{c}_{III}(a, e) + \sum_{b \neq 0, c \neq 0} \mu_{bc}^{(1)} \mathbf{c}_{IV}(b, c) \\ & + \sum_{b \neq 0, f} \nu_{bf}^{(4)} \mathbf{c}_V(b, f) + \sum_{c \neq 0, d \neq 0} \mu_{cd}^{(2)} \mathbf{c}_{VI}(c, d) + \sum_{c \neq 0} \nu_c^{(2)} \mathbf{c}_{VII}(c) \\ & + \sum_{d \neq 0, e} \lambda_{de}^{(1)} \mathbf{c}_{VIII}(d, e) + \sum_{e \neq 0} \sigma_e^{(2)} \mathbf{c}_{IX}(e) + \sum_f \nu_f^{(1)} \mathbf{c}_X(f) \end{aligned}$$

satisfies $\theta^+(y) = (y_0, y_1, \tilde{y}_2, y_2, y_3)$.

Therefore θ^+ induces an isomorphism between $\mathbb{Z}_p[\text{Conj}(G^{(n)})]$ and $\Omega^{(n)}$. $\square$

5. TRANSLATION INTO THE MULTIPLICATIVE THETA MAP

In the previous section, we construct the additive theta map θ^+ . Now we shall translate it into *the multiplicative theta map* θ . The main tool for this translation is the integral logarithmic homomorphism introduced in §1.3.

Let $\theta_i: K_1(\Lambda(G)) \rightarrow \Lambda(U_i/V_i)^\times$ be the composition of the norm homomorphism of K -groups

$$\text{Nr}_i: K_1(\Lambda(G)) \rightarrow K_1(\Lambda(U_i))$$

and the natural homomorphism $K_1(\Lambda(U_i)) \rightarrow K_1(\Lambda(U_i/V_i)) = \Lambda(U_i/V_i)^\times$ induced by $\Lambda(U_i) \rightarrow \Lambda(U_i/V_i)$. Set

$$\theta = (\theta_i)_i: K_1(\Lambda(G)) \rightarrow \prod_i \Lambda(U_i/V_i)^\times.$$

Proposition-Definition 5.1 (The Frobenius homomorphism). *For an arbitrary element $g \in G$, set*

$$(5.1) \quad \varphi(g) = g^p.$$

Then φ induces a group homomorphism $\varphi: G \rightarrow \Gamma$. We call this homomorphism the Frobenius homomorphism.

We denote the induced ring homomorphism $\Lambda(G) \rightarrow \Lambda(\Gamma)$ by the same symbol φ , and also call it *the Frobenius homomorphism*.

Proof. Since the exponent of G^f is exactly equal to p , we may decompose φ as follows:

$$\begin{array}{ccc}
G & \xrightarrow{\varphi} & \Gamma \\
\pi_G \searrow & & \nearrow \tilde{\varphi} \\
& \Gamma &
\end{array}$$

where $\tilde{\varphi}: \Gamma \rightarrow \Gamma$ is the Frobenius endomorphism of Γ defined by $t \mapsto t^p$, and π_G is the canonical surjection. Hence φ is clearly a group homomorphism. $\square$

Remark 5.2. In general, the correspondence (5.1) does not induce a group homomorphism $\varphi: G \rightarrow G$.

In the following, we denote by the same symbol φ the induced ring homomorphism $\Lambda(U_i/V_i) \rightarrow \Lambda(\Gamma)$.

Definition 5.3. We define Ψ to be the subgroup of $\prod_i \Lambda(U_i/V_i)^\times$ consisting of all elements $(\eta_0, \eta_1, \tilde{\eta}_2, \eta_2, \eta_3)$ satisfying the following two conditions:

(1) (norm relations)

$$\text{(rel-1)} \quad \text{Nr}_{\Lambda(U_0/V_0)/\Lambda(U_1/V_0)} \eta_0 \equiv \eta_1,$$

$$\text{(rel-2)} \quad \text{Nr}_{\Lambda(U_0/V_0)/\Lambda(\tilde{U}_2/V_0)} \eta_0 \equiv \tilde{\eta}_2,$$

$$\text{(rel-3)} \quad \text{Nr}_{\Lambda(\tilde{U}_2/\tilde{V}_2)/\Lambda(U_2/\tilde{V}_2)} \tilde{\eta}_2 \equiv \eta_2,$$

$$\text{(rel-4)} \quad \text{Nr}_{\Lambda(U_1/\tilde{V}_2)/\Lambda(U_1 \cap \tilde{U}_2/\tilde{V}_2)} \eta_1 \equiv \text{Nr}_{\Lambda(\tilde{U}_2/\tilde{V}_2)/\Lambda(U_1 \cap \tilde{U}_2/\tilde{V}_2)} \tilde{\eta}_2,$$

$$\text{(rel-5)} \quad \text{Nr}_{\Lambda(U_1/V_1)/\Lambda(U_3/V_1)} \eta_1 \equiv \eta_3,$$

$$\text{(rel-6)} \quad \text{Nr}_{\Lambda(U_2/V_2)/\Lambda(U_3/V_2)} \eta_2 \equiv \eta_3,$$

(See Figure 1).

(2) (congruences)

$$\eta_1 \equiv \varphi(\eta_0) \pmod{I_1}, \quad \tilde{\eta}_2 \equiv \varphi(\eta_0) \pmod{\tilde{I}_2},$$

$$\eta_2 \equiv \varphi(\eta_0)^p \pmod{I_2}, \quad \eta_3 \equiv \varphi(\eta_0)^{p^2} \pmod{I_3}.$$

Now let us recall the definition of norm maps of commutative rings: let R be a commutative ring and R' a commutative R -algebra. Assume that R' is free and finitely generated as an R -module. Then we define $\text{Nr}_{R'/R}(y)$ to be the determinant of the multiplication- y homomorphism.

Proposition 5.4. *The homomorphism θ induces a surjection*

$$(5.2) \quad \theta: K_1(\Lambda(G)) \rightarrow \Psi.$$

In other words, θ induces the (multiplicative) theta map for G (in the sense of Definition 3.3).

In the rest of this section, we prove Proposition 5.4 by using the additive theta map θ^+ and the integral logarithmic homomorphisms. Since the integral logarithmic homomorphisms are defined only for group rings of finite quotients, we fix $n \geq 1$ and construct an isomorphism for finite quotients

$$\theta^{(n)}: K_1(\mathbb{Z}_p[G^{(n)}])/SK_1(\mathbb{Z}_p[G^{(n)}]) \xrightarrow{\sim} \Psi^{(n)} \left(\subseteq \prod_i \mathbb{Z}_p[U_i^{(n)}/V_i] \right)$$

where $G^{(n)} = G_f \times \Gamma/\Gamma^{p^n}$, $U_i^{(n)} = U_i^f \times \Gamma/\Gamma^{p^n}$, and $\Psi^{(n)}$ is the subgroup of $\prod_i \mathbb{Z}_p[U_i^{(n)}/V_i]$ defined by the same conditions of Definition 5.3. Then we obtain the surjection (5.2) by taking the projective limit.

To simplify the notation, we denote the theta map for $G^{(n)}$ by θ in §5.1, §5.2, and §5.3. In §5.4, where we take the projective limit of theta maps, we denote the theta map for $G^{(n)}$ by $\theta^{(n)}$ again.

5.1. Logarithmic isomorphisms. In the following three subsections, we fix $n \geq 1$.

For each $i \geq 1$, we define the ideal $J_i^{(n)}$ of $\mathbb{Z}_p[U_i^{(n)}/V_i]$ as follows:

$$\begin{aligned} J_1^{(n)} &= (h_{\varepsilon^j \delta^k} \quad (0 \leq j, k \leq p-1)), & \tilde{J}_2^{(n)} &= (p, h_\delta), \\ J_2^{(n)} &= J_3^{(n)} = (p, h_\zeta). \end{aligned}$$

Note that $J_i^{(n)}$ contains $I_i^{(n)}$. Since $h_{\varepsilon^0 \delta^0} = p$, $J_1^{(n)}$ also contains p .

Lemma 5.5. *For each $i \geq 1$, $1 + J_i^{(n)}$ is a multiplicative group and the p -adic logarithmic homomorphism induces an isomorphism $1 + J_i^{(n)} \xrightarrow{\sim} J_i^{(n)}$.*

Proof. Since $J_i^{(n)}$ is an ideal, $(J_i^{(n)})^2 \subseteq J_i^{(n)}$. Therefore $1 + J_i^{(n)}$ is closed under multiplication.

It is easy to calculate that $h_\delta^2 = ph_\delta$ and $h_\zeta^2 = ph_\zeta$. Then for $\tilde{J}_2^{(n)}$, $J_2^{(n)}$ and $J_3^{(n)}$, we have $(J_i^{(n)})^2 = pJ_i^{(n)}$. For $J_1^{(n)}$, we have

$$h_{\varepsilon^j \delta^k} h_{\varepsilon^{j'} \delta^{k'}} = \begin{cases} h_{\varepsilon, \delta} & \text{if } (j, k) \text{ and } (j', k') \text{ are } \mathbb{Z}\text{-linearly independent,} \\ ph_{\varepsilon^j \delta^k} & \text{otherwise,} \end{cases}$$

where

$$h_{\varepsilon, \delta} = \sum_{0 \leq \ell, \ell' \leq p-1} \varepsilon^\ell \delta^{\ell'}.$$

For each $0 \leq j, k \leq p-1$, we have $h_{\varepsilon, \delta} h_{\varepsilon^j \delta^k} = ph_{\varepsilon, \delta}$ and $h_{\varepsilon, \delta}^2 = p^2 h_{\varepsilon, \delta}$ (note that the exponent of U_1^f/V_1^f is p). Thus we have

$$(J_1^{(n)})^3 = p(J_1^{(n)})^2 \subseteq pJ_1^{(n)}.$$

Set $N = 3$ for $i = 1$ and $N = 2$ for other i .

- The existence of inverse elements.

By the argument above, we have $y^m \in p^{m-N+1} J_i^{(n)}$ for an arbitrary element $y \in J_i^{(n)}$ and every $m \geq N$. Therefore,

$$(5.3) \quad (1 + y)^{-1} = \sum_{m \geq 0} (-1)^m y^m$$

converges in $J_i^{(n)}$ p -adically.

- Convergence of logarithms.

It is obvious that $y^m/m^i n J_i^{(n)}$ for $1 \leq m < N$, and we have

$$\frac{y^m}{m} \in \frac{p^{m-N+1}}{m} \cdot J_i^{(n)} \subseteq J_i^{(n)}$$

for an arbitrary $y \in J_i^{(n)}$ and every $m \geq N$ (here we use the fact that $p \neq 0$). Thus the logarithm $\log(1+y) = \sum_{m \geq 1} (-1)^{m-1} (y^m/m)$ converges p -adically in $J_i^{(n)}$.

Since $\left\{ (J_i^{(n)})^m \right\}^N = p^m (J_i^{(n)})^m (N-1) \subseteq p^m (J_i^{(n)})^m$, we may also show that $1 + (J_i^{(n)})^m$ is a subgroup of $1 + J_i^{(n)}$ and

$$\log \left(1 + (J_i^{(n)})^m \right) \subseteq (J_i^{(n)})^m \quad \text{for every } m \geq 1,$$

by the same argument as above.

- Logarithmic isomorphisms.

First note that p -adic topology and $J_i^{(n)}$ -adic topology are the same on $J_i^{(n)}$: this is because $p^{m+1} J_i^{(n)} \subseteq (J_i^{(n)})^{m+N-1} \subseteq p^m J_i^{(n)}$ for every $m \geq 1$. Since $J_i^{(n)}$ is p -adically complete, it is also $J_i^{(n)}$ -adically complete. Therefore to show that the logarithm induces an isomorphism $1 + J_i^{(n)} \xrightarrow{\cong} J_i^{(n)}$, it is sufficient to show that it induces an isomorphism

$$\begin{aligned} \log: (1 + (J_i^{(n)})^m) / (1 + (J_i^{(n)})^{m+1}) &\xrightarrow{\cong} (J_i^{(n)})^m / (J_i^{(n)})^{m+1} \\ &; 1 + y \mapsto y \end{aligned}$$

for each $m \geq 1$, and to prove this, it suffices to show that

$$(5.4) \quad y^{p^k} / p^k \in (J_i^{(n)})^{m+1} \text{ for every } 1 + y \in 1 + (J_i^{(n)})^m \text{ and } k \geq 1.$$

However, since $y^{N+1} \in p (J_i^{(n)})^{m+1}$, we obtain $y^p \in p (J_i^{(n)})^{m+1}$ (recall $p \neq 2, 3$ by assumption). This implies (5.4). $\square$

Lemma 5.6. *For each $i \geq 1$, $1 + I_i^{(n)}$ is a multiplicative group and the p -adic logarithmic homomorphism induces an isomorphism $1 + I_i^{(n)} \xrightarrow{\cong} I_i^{(n)}$.*

Remark 5.7. Since each $I_i^{(n)}$ is **not** an ideal but only a $\mathbb{Z}_p$ -submodule of $\mathbb{Z}_p[U_i^{(n)}/V_i]$, it is not trivial even that $1 + I_i^{(n)}$ is closed under multiplication.

Proof. By direct calculation, we have

$$\begin{aligned} (I_1^{(n)})^2 &= [p\alpha^a \gamma^c \delta^d \varepsilon^e h_{\varepsilon^c \delta^{-a}}, p\delta^d \varepsilon^e h_{\varepsilon^c \delta^{-a}}, \alpha^a \gamma^c h_{\varepsilon, \delta}((a, c) \neq (0, 0))]_{\mathbb{Z}_p[\Gamma/\Gamma p^n]}, \\ (\tilde{I}_2^{(n)})^2 &= [p\beta^b \gamma^c h_\delta, p^2 \gamma^c \delta^d]_{\mathbb{Z}_p[\Gamma/\Gamma p^n]}, \\ (I_2^{(n)})^2 &= [p\beta^b \gamma^c h_\zeta, p^2 \beta^b \zeta^f]_{\mathbb{Z}_p[\Gamma/\Gamma p^n]}, \\ (I_3^{(n)})^2 &= [p^6 \zeta^f, p^5 \varepsilon^e h_\zeta, p^4 \gamma^c h_\varepsilon h_\zeta]_{\mathbb{Z}_p[\Gamma/\Gamma p^n]}, \end{aligned}$$

These results imply that $(I_i^{(n)})^2 \subseteq I_i^{(n)}$.⁶ Therefore, each $1 + I_i^{(n)}$ is stable under multiplication.

⁶For $I_1^{(n)}$, recall that $I_1^{(n)}$ is also stable under the multiplication of δ and ε . Therefore we have $p\delta^d \varepsilon^e h_{\varepsilon^c \delta^{-a}} = \left(\sum_{\ell=0}^{p-1} \delta^{d-\ell a} \varepsilon^{e+\ell c} \right) p \in I_1^{(n)}$ and $\alpha^a \gamma^c h_{\varepsilon, \delta} =$

Now let

$$\begin{aligned} I'_1 &= I_1^{(n)}, \\ \tilde{I}'_2 &= [\beta^b \gamma^c h_\delta, p\gamma^c \delta^d]_{\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]}, \\ I'_2 &= [p\beta^b \zeta^f, \beta^b \gamma^c h_\zeta]_{\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]}, \\ I'_3 &= [p^3 \zeta^f, p^2 \varepsilon^e h_\zeta, p\gamma^c h_\varepsilon h_\zeta]_{\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]}. \end{aligned}$$

Note that $I'_i \supseteq I_i^{(n)}$, and by simple calculation we have $(I_1^{(n)})^3 = (I'_1)^3 \subseteq p(I'_1)^2$ for $i = 1$ and $(I_i^{(n)})^2 = (I'_i)^2 \subseteq pI'_i$ for other i .

Then we may show the existence of inverse elements of $1 + I'_i$ and the logarithmic isomorphism $\log: 1 + I'_i \xrightarrow{\sim} I'_i$ for each i by the same argument as the proof of Lemma 5.5 by replacing J_i by I'_i . Since $(I'_i)^2 \subseteq I_i^{(n)} \subseteq I'_i$ and $I_i^{(n)}$ is a closed subset of I'_i , the power series $(1 + x)^{-1}$ and the p -adic logarithm $\log(1 + x)$ on $1 + I_i^{(n)}$ converges into $I_i^{(n)}$. Moreover since $\log: 1 + (I'_i)^k \rightarrow (I'_i)^k$ is an isomorphism for $k = 1, 2$ and

$$\log: 1 + I_i^{(n)} / 1 + (I'_i)^2 \longrightarrow I_i^{(n)} / (I'_i)^2; 1 + y_i \mapsto y_i \pmod{(I'_i)^2}$$

is also an isomorphism, we may conclude that $\log: 1 + I_i^{(n)} \rightarrow I_i^{(n)}$ is an isomorphism. $\square$

5.2. $\Psi^{(n)}$ contains the image of θ .

Lemma 5.8. *The following diagram commutes for each i and $n \geq 1$:*

$$\begin{array}{ccc} K_1(\mathbb{Z}_p[G^{(n)}]) & \xrightarrow{\log} & \mathbb{Z}_p[\text{Conj}(G^{(n)})] \\ \text{Nr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]} \downarrow & & \downarrow \text{Tr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]} \\ K_1(\mathbb{Z}_p[U_i^{(n)}]) & \xrightarrow{\log} & \mathbb{Z}_p[\text{Conj}(U_i^{(n)})] \end{array}$$

Proof. By the proof of [Oli-Tay], Theorem 1.4., the following diagram commutes.

$$\begin{array}{ccc} K_1(\mathbb{Z}_p[G^{(n)}]) & \xrightarrow{\log} & \mathbb{Z}_p[\text{Conj}(G^{(n)})] \\ \text{Nr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]} \downarrow & & \downarrow R' \\ K_1(\mathbb{Z}_p[U_i^{(n)}]) & \xrightarrow{\log} & \mathbb{Z}_p[\text{Conj}(U_i^{(n)})] \end{array}$$

where $R': \mathbb{Z}_p[\text{Conj}(G^{(n)})] \rightarrow \mathbb{Z}_p[\text{Conj}(U_i^{(n)})]$ is defined as follows: for an arbitrary $x \in G^{(n)}$, let $\{u'_1, \dots, u'_{s_i}\}$ be a set of representatives of the double coset decomposition $\langle x \rangle \backslash G^{(n)} / U_i^{(n)}$, and let

$$J = \{j \mid 1 \leq j \leq s_i, u_j'^{-1} x u'_j \in U_i^{(n)}\}.$$

Then we define $R'(x) = \sum_{j \in J} u_j'^{-1} x u'_j$.

$\overline{\left(\sum_{\ell=0}^{p-1} (\varepsilon^{c'} \delta^{-a'})^\ell \right)} \alpha^a \gamma^c h_{\varepsilon^c \delta^{-a}} \in I_1^{(n)}$ where $(a', c') \in \mathbb{Z}^2$ is an arbitrary element $\mathbb{Z}$ -linearly independent of (a, c) . Therefore we have $(I_1^{(n)})^2 \subseteq I_1^{(n)}$.

It suffices to show that

$$(5.5) \quad R' = \text{Tr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]},$$

but this is not difficult at all.

Let $\{u'_1, \dots, u'_{s_i}\}$ be as above. Then we can write

$$G^{(n)} = \bigsqcup_j \langle x \rangle u'_j U_i^{(n)} = \bigcup_{k=0}^{p-1} \bigsqcup_j x^k u'_j U_i^{(n)},$$

therefore it is clear that $\{x^k u'_j\}_{0 \leq k \leq p-1, j}$ contains a set of representatives of the right coset decomposition $G^{(n)}/U_i^{(n)}$. Moreover, since each $u'_j U_i^{(n)}$ is disjoint, $\{u'_j\}_j$ is a subset of representatives of $G^{(n)}/U_i^{(n)}$.

$$(\text{Case-1}) \quad G^{(n)} = \bigsqcup_j u'_j U_i^{(n)}.$$

In this case, the desired equation (5.5) is obvious by definition of R' and Tr .

$$(\text{Case-2}) \quad G^{(n)} = \bigcup_{k=0}^{p-1} \bigsqcup_j x^k u'_j U_i^{(n)}.$$

Then by simple calculation

$$\begin{aligned} R'(x) &= \sum_{j \in J} u'_j{}^{-1} x u'_j = \frac{1}{p} \sum_{k=0}^{p-1} \sum_{j \in J} (x^k u'_j)^{-1} x (x^k u'_j) \\ &= \frac{1}{p} \text{Tr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]} x. \end{aligned}$$

On the other hand, recall that $j \in J$ implies that $x u'_j$ is an element of $u'_j U_i^{(n)}$. However, this is impossible since $\{x^k u'_j\}_{0 \leq k \leq p-1, j}$ is a set of representatives of $G^{(n)}/U_i^{(n)}$ in this case. Therefore $J = \emptyset$ and $R'(x) = \text{Tr}_{\mathbb{Z}_p[G^{(n)}]/\mathbb{Z}_p[U_i^{(n)}]}(x) = 0$.

□

Proposition 5.9. *For an arbitrary element $\eta \in K_1(\mathbb{Z}_p[G^{(n)}])$, the following equations hold:*

$$\begin{aligned} \theta_1^+ \circ \Gamma_{G^{(n)}}(g) &= \log \frac{\theta_1(\eta)}{\varphi(\theta_0(\eta))}, \\ \tilde{\theta}_2^+ \circ \Gamma_{G^{(n)}}(g) &= \log \frac{\tilde{\theta}_2(\eta)}{\varphi(\theta_0(\eta))}, \\ \theta_2^+ \circ \Gamma_{G^{(n)}}(g) &= \log \frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p}, \\ \theta_3^+ \circ \Gamma_{G^{(n)}}(g) &= \log \frac{\theta_3(\eta)}{\varphi(\theta_0(\eta))^{p^2}}, \end{aligned}$$

where $\Gamma_{G^{(n)}}: K_1(\mathbb{Z}_p[G^{(n)}]) \rightarrow \mathbb{Z}_p[\text{Conj}(G^{(n)})]$ is the integral logarithm for $G^{(n)}$ (Proposition-Definition 1.29).

Proof. This proposition follows from Lemma 5.8 and by following commutative diagrams:

$$\begin{array}{ccc}
\mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_0^+} & \mathbb{Z}_p[U_0^{(n)}/V_0] & \mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_0^+} & \mathbb{Z}_p[U_0^{(n)}/V_0] \\
\frac{1}{p}\varphi \downarrow & & \downarrow \varphi & \frac{1}{p}\varphi \downarrow & & \downarrow \varphi \\
\mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_1^+} & \mathbb{Z}_p[U_1^{(n)}/V_1] & \mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\tilde{\theta}_2^+} & \mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2] \\
\\
\mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_0^+} & \mathbb{Z}_p[U_0^{(n)}/V_0] & \mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_0^+} & \mathbb{Z}_p[U_0^{(n)}/V_0] \\
\frac{1}{p}\varphi \downarrow & & \downarrow p\varphi & \frac{1}{p}\varphi \downarrow & & \downarrow p^2\varphi \\
\mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_2^+} & \mathbb{Z}_p[U_2^{(n)}/V_2] & \mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\theta_3^+} & \mathbb{Z}_p[U_3^{(n)}/V_3]
\end{array}$$

These diagrams are easily checked for an arbitrary element of $G^{(n)}$. Then we may calculate as

$$\begin{aligned}
\theta_2^+ \circ \Gamma_{G^{(n)}}(\eta) &= \theta_2^+ \left(\log \eta - \frac{1}{p} \varphi(\log(\eta)) \right) \\
&= \theta_2^+(\log(\eta)) - p\varphi(\theta_0^+(\log(\eta))) \\
&= \log(\theta_2(\eta)) - p\varphi(\log(\theta_0(\eta))) \\
&= \log \frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p},
\end{aligned}$$

here we use the diagram above for the second equality, and use Lemma 5.8 for the third equality. The other equations may be derived similarly.

Note that for an arbitrary $y \in \mathbb{Z}_p[U_i^{(n)}/V_i]$, we have

$$\varphi(\log(y)) = \log \varphi(y)$$

by the definition of the p -adic logarithms and the fact that the Frobenius homomorphism φ is a ring homomorphism on each $\mathbb{Z}_p[U_i^{(n)}/V_i]$. $\square$

Lemma 5.10. *For an arbitrary element $\eta \in K_1(\mathbb{Z}_p[G^{(n)}])$, the following congruences hold:*

$$\begin{aligned}
\theta_1(\eta) &\equiv \varphi(\theta_0(\eta)) && \text{mod } J_1^{(n)}, \\
\tilde{\theta}_2(\eta) &\equiv \varphi(\theta_0(\eta)) && \text{mod } \tilde{J}_2^{(n)}, \\
\theta_2(\eta) &\equiv \varphi(\theta_0(\eta))^p && \text{mod } J_2^{(n)}, \\
\theta_3(\eta) &\equiv \varphi(\theta_0(\eta))^{p^2} && \text{mod } J_3^{(n)}.
\end{aligned}$$

Proof. We calculate the image of the theta map as in §1.2.

In the following proof, we use the same notation η for a lift of η to $\mathbb{Z}_p[G^{(n)}]^\times$ (Proposition 1.7).

- The congruences for $\theta_1, \tilde{\theta}_2$.

We only show the congruence for $\tilde{\theta}_2$. We may prove the congruence for θ_1 in the same manner.

Since $\{\alpha^i \mid 0 \leq i \leq p-1\}$ is a $\mathbb{Z}_p[\widetilde{U}_2^{(n)}]$ -basis of $\mathbb{Z}_p[G^{(n)}]$, η is described as a $\mathbb{Z}_p[\widetilde{U}_2^{(n)}]$ -linear combination as follows:

$$\eta = \sum_{i=0}^{p-1} \eta_i \alpha^i, \quad \eta_i \in \mathbb{Z}_p[\widetilde{U}_2^{(n)}].$$

Then we have

$$\begin{aligned} \alpha^j \eta &= \sum_{i=0}^{p-1} (\alpha^j \eta_i \alpha^{-j}) \cdot \alpha^{i+j} \\ &= \sum_{i=0}^{p-1} \nu_j(\eta_{i-j}) \alpha^i \end{aligned}$$

where $\nu_j: \mathbb{Z}_p[\widetilde{U}_2^{(n)}] \rightarrow \mathbb{Z}_p[\widetilde{U}_2^{(n)}]; x \mapsto \alpha^j x \alpha^{-j}$. Here we consider the sub-index of η_i as an element of $\mathbb{Z}/p\mathbb{Z}$. By abuse of notation, we denote the image of η_i in $\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$ by the same symbol η_i . Then we may compute $\widetilde{\theta}_2(\eta)$ as follows:

$$\begin{aligned} \widetilde{\theta}_2(\eta) &= \det(\nu_j(\eta_{i-j}))_{i,j} \\ &= \sum_{\sigma \in \mathfrak{S}_p} \text{sgn}(\sigma) \prod_{j=0}^{p-1} \nu_j(\eta_{\sigma(j)-j}) \\ &= \sum_{\sigma \in \mathfrak{S}_p} P_\sigma \end{aligned}$$

where $\mathfrak{S}_p$ is the permutation group of $\{0, 1, \dots, p-1\}$ and $P_\sigma = \text{sgn}(\sigma) \prod_{j=0}^{p-1} \nu_j(\eta_{\sigma(j)-j})$.

Now we have

$$\begin{aligned} \nu_k(P_\sigma) &= \text{sgn}(\sigma) \prod_j \nu_{j+k}(\eta_{\sigma(j)-j}) \\ &= \text{sgn}(\sigma) \prod_j \nu_j(\eta_{(\sigma(j-k)+k)-j}). \end{aligned}$$

First Suppose that σ does not satisfy

$$(5.6) \quad \sigma(j-k) + k = \sigma(j) \quad \text{for each } k \in \mathbb{Z}/p\mathbb{Z},$$

then

$$\tau_k(j) = \sigma(j-k) + k, \quad k \in \mathbb{Z}/p\mathbb{Z}$$

are distinct elements of $\mathfrak{S}_p$ and we have

$$(5.7) \quad \nu_k(P_\sigma) = P_{\tau_k}.$$

Here we use the fact $\text{sgn}(\sigma) = \text{sgn}(\tau_k)$. This follows from the equation $\tau_k = c_k \circ \sigma \circ c_k^{-1}$ where $c_k(i) = i + k$.

The equation (5.7) implies

$$\sum_{k=0}^{p-1} P_{\tau_k} \in \sum_{k=0}^{p-1} \nu_k \left(\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2] \right).$$

Note that each ν_k is a $\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$ -linear map where

$$\widetilde{U}_2^{(n)} = \langle \gamma, \delta, \varepsilon, \zeta \rangle \times \Gamma/\Gamma^{p^n}$$

and $\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$ is generated by $\{\beta^b \mid b \in \mathbb{Z}/p\mathbb{Z}\}$ over $\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$. Since

$$\sum_{k=0}^{p-1} \nu_k(\beta^b) = \begin{cases} p & \text{if } b = 0, \\ \beta^b(1 + \delta + \dots + \delta^{p-1}) & \text{otherwise,} \end{cases}$$

we may conclude that

$$(5.8) \quad \sum_{k=0}^{p-1} \nu_k \left(\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2] \right) \subseteq \widetilde{J}_2^{(n)}.$$

This implies $\sum_{k=0}^{p-1} P_{\tau_k} \in \widetilde{J}_2^{(n)}$.

Next assume σ satisfies (5.6). It is easy to see that all permutations satisfying (5.6) are

$$c_h(j) = j + h \quad (0 \leq h \leq p-1).$$

For $\sigma = c_h$, we have

$$P_{c_h} = \text{sgn}(c_h) \prod_{i=0}^{p-1} \nu_j(\eta_{c_h(j)-j}) = \prod_{i=0}^{p-1} \nu_j(\eta_h).$$

Note that since c_h ($h \neq 0$) is a cyclic permutation of degree p , c_h is an even permutation, therefore $\text{sgn}(c_h) = 1$ for $0 \leq h \leq p-1$.

Claim. For an arbitrary element $x \in \mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$,

$$\prod_{j=0}^{p-1} \nu_j(x) \equiv \varphi(x) \pmod{\widetilde{J}_2^{(n)}}.$$

If this claim is true, we have

$$\begin{aligned} \widetilde{\theta}_2(\eta) &\equiv \sum_{h=0}^{p-1} P_{c_h} && \pmod{\widetilde{J}_2^{(n)}} \\ &\equiv \sum_{h=0}^{p-1} \prod_{j=0}^{p-1} \nu_j(\eta_h) && \pmod{\widetilde{J}_2^{(n)}} \\ &\equiv \sum_{h=0}^{p-1} \varphi(\eta_h) && \pmod{\widetilde{J}_2^{(n)}} \\ &\equiv \sum_{h=0}^{p-1} \varphi(\eta_h \beta^h) && \pmod{\widetilde{J}_2^{(n)}} \\ &\equiv \varphi(\eta) && \pmod{\widetilde{J}_2^{(n)}}, \end{aligned}$$

and the congruence for $\widetilde{\theta}_2$ holds.

Now let

$$x = \sum_{\ell=1}^N x_\ell$$

where each x_ℓ is a monomial of $\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2]$. Then

$$\begin{aligned}
 \prod_{j=0}^{p-1} \nu_j(x) &= \prod_{j=0}^{p-1} \sum_{\ell=1}^N \nu_j(x_\ell) \\
 (5.9) \quad &= \sum_{1 \leq \ell_0, \dots, \ell_{p-1} \leq N} \nu_0(x_{\ell_0}) \nu_1(x_{\ell_1}) \cdots \nu_{p-1}(x_{\ell_{p-1}}) \\
 &= \sum_{1 \leq \ell_0, \dots, \ell_{p-1} \leq N} Q_{\ell_0, \dots, \ell_{p-1}}.
 \end{aligned}$$

Here we set $Q_{\ell_0, \dots, \ell_{p-1}} = \nu_0(x_{\ell_0}) \nu_1(x_{\ell_1}) \cdots \nu_{p-1}(x_{\ell_{p-1}})$.

If $\ell_0 = \ell_1 = \dots = \ell_{p-1} = \lambda$ (constant), we obtain $Q_{\lambda, \dots, \lambda} = \varphi(x_\lambda)$ by easy calculation (here we use that the exponent of G^f is p). Otherwise, each

$$Q_{\ell_k, \ell_{k+1}, \dots, \ell_{p-1}, \ell_0, \ell_1, \dots, \ell_{k-1}} = \nu_{p-k}(Q_{\ell_0, \dots, \ell_{p-1}})$$

is a distinct term in the expansion (5.9), so we have

$$\sum_{k=0}^{p-1} Q_{\ell_k, \ell_{k+1}, \dots, \ell_{p-1}, \ell_0, \ell_1, \dots, \ell_{k-1}} \in \sum_{k=0}^{p-1} \nu_k \left(\mathbb{Z}_p[\widetilde{U}_2^{(n)}/\widetilde{V}_2] \right),$$

which is contained in $\widetilde{J}_2^{(n)}$ by (5.8). Hence,

$$\begin{aligned}
 \prod_{j=0}^{p-1} \nu_j(x) &\equiv \sum_{\lambda=1}^N \varphi(x_\lambda) \pmod{\widetilde{J}_2^{(n)}} \\
 &\equiv \varphi(x) \pmod{\widetilde{J}_2^{(n)}},
 \end{aligned}$$

and Claim is proven.

- The congruence for θ_2 .

The basic strategy is essentially the same as the congruence for $\widetilde{\theta}_2$, but since U_2 is not a normal subgroup of G , the calculation is much more complicated.

Since $\{\alpha^i \delta^j \mid 0 \leq i, j \leq p-1\}$ is a system of representatives of $U_2^{(n)} \backslash G^{(n)}$ and $\{\beta^k \mid 0 \leq k \leq p-1\}$ is that of $U_3^{(n)} \backslash U_2^{(n)}$, we can describe η in the following form:

$$\eta = \sum_{0 \leq i, j, k \leq p-1} (\eta_{i,j}^{(k)} \beta^k) \alpha^i \delta^j$$

where $\eta_{i,j}^{(k)}$ is an element of $\mathbb{Z}_p[U_3^{(n)}]$. Then we have

$$\begin{aligned}\alpha^\ell \delta^m \eta &= \sum_{0 \leq i,j,k \leq p-1} \nu_{\ell,m}(\eta_{i,j}^{(k)} \beta^k) \alpha^{i+\ell} \delta^{j+m} \\ &= \sum_{0 \leq i,j,k \leq p-1} \{\nu_{\ell,m}(\eta_{i,j}^{(k)}) \delta^{\ell k} \beta^k\} \alpha^{i+\ell} \delta^{j+m} \\ &= \sum_{0 \leq i,j \leq p-1} \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{i-\ell,j-m-\ell k}^{(k)}) \beta^k \right) \alpha^i \delta^j\end{aligned}$$

where $\nu_{\ell,m}(x) = (\alpha^\ell \delta^m) x (\alpha^\ell \delta^m)^{-1}$. We use the fundamental relations $[\alpha, \delta] = 1$ and $[\alpha, \beta] = \delta$ for the second equality.

Here we regard the sub-index i, j of $\eta_{i,j}^{(k)}$ as an element of $(\mathbb{Z}/p\mathbb{Z})^{\oplus 2}$, and the upper-index (k) of it as an element of $\mathbb{Z}/p\mathbb{Z}$. We denote the image of $\eta_{i,j}^{(k)}$ in $\mathbb{Z}_p[U_3^{(n)}/V_2^{(n)}]$ by the same symbol $\eta_{i,j}^{(k)}$. Then we obtain

$$\begin{aligned}\theta_2(\eta) &= \det \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{(i,j)-(\ell,m)-(0,\ell k)}^{(k)}) \beta^k \right)_{(i,j),(\ell,m)} \\ &= \sum_{\sigma \in \mathfrak{S}_\square} \text{sgn}(\sigma) \prod_{(\ell,m) \in (\mathbb{Z}/p\mathbb{Z})^{\oplus 2}} \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{\sigma(\ell,m)-(\ell,m)-(0,\ell k)}^{(k)}) \beta^k \right) \\ &= \sum_{\sigma \in \mathfrak{S}_\square} P_\sigma\end{aligned}$$

where $\mathfrak{S}_\square$ is the permutation group of $\{(\ell, m) \mid 0 \leq \ell, m \leq p-1\}$.

Now by using $[\beta, \delta] = 1$, we have

$$\begin{aligned}&\nu_{0,\mu}(P_\sigma) \\ &= \text{sgn}(\sigma) \prod_{0 \leq \ell, m \leq p-1} \left(\sum_{k=0}^{p-1} \nu_{\ell,m+\mu}(\eta_{\sigma(\ell,m)-(\ell,m)-(0,\ell k)}^{(k)}) \beta^k \right) \\ &= \text{sgn}(\sigma) \prod_{0 \leq \ell, m \leq p-1} \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{(\sigma(\ell,m-\mu)+(0,\mu))-(\ell,m)-(0,\ell k)}^{(k)}) \beta^k \right).\end{aligned}$$

Therefore if σ does not satisfy

$$(5.10) \quad \sigma(\ell, m) = \sigma(\ell, m - \mu) + (0, \mu) \quad \text{for each } \mu \in \mathbb{Z}/p\mathbb{Z},$$

then

$$\tau_\mu(\ell, m) = \sigma(\ell, m - \mu) + (0, \mu), \quad \mu \in \mathbb{Z}/p\mathbb{Z}$$

are distinct elements of $\mathfrak{S}_\square$, and we obtain

$$\sum_{\mu=0}^{p-1} P_{\tau_\mu} \in \sum_{\mu} \nu_{0,\mu} \left(\mathbb{Z}_p[U_2^{(n)}/V_2] \right),$$

here we use the fact $\text{sgn}(\sigma) = \text{sgn}(\tau_\mu)$ which can be proven by the same argument as the case of $\tilde{\theta}_2$.

Note that each $\nu_{0,\mu}$ is a $\mathbb{Z}_p[U_2'^{(n)}/V_2]$ -linear map where

$$U_2'^{(n)} = \langle \beta, \varepsilon, \zeta \rangle \times \Gamma/\Gamma^{p^n}$$

and $\mathbb{Z}_p[U_2'^{(n)}/V_2]$ is generated by $\{\gamma^c \mid c \in \mathbb{Z}/p\mathbb{Z}\}$ over $\mathbb{Z}_p[U_2'^{(n)}/V_2]$. Since

$$\sum_{\mu=0}^{p-1} \nu_{0,\mu}(\gamma^c) = \begin{cases} p & \text{if } c = 0, \\ \gamma^c(1 + \zeta + \cdots + \zeta^{p-1}) & \text{otherwise,} \end{cases}$$

we may conclude that

$$(5.11) \quad \sum_{\mu=0}^{p-1} \nu_{0,\mu} \left(\mathbb{Z}_p[U_2'^{(n)}/V_2] \right) \subseteq J_2^{(n)}$$

and this implies $\sum_{k=0}^{p-1} P_{\tau_k} \in \tilde{J}_2^{(n)}$ (note that the argument up to here is almost the same as the case of $\tilde{\theta}_2$).

Now suppose that $\sigma \in \mathfrak{S}_{\square}$ satisfies (5.10). If we set $\sigma(\ell, 0) = (a_\ell, b_\ell)$, then by (5.10) we have

$$\sigma(\ell, -\mu) = (a_\ell, b_\ell - \mu) \quad \text{for each } \mu \in \mathbb{Z}/p\mathbb{Z}.$$

This calculation implies that all permutations satisfying (5.10) are described in the following forms:

$$(5.12) \quad c_{s,h}(\ell, m) = (s(\ell), h_\ell + m)$$

where s is a permutation of $\{0, 1, \dots, p-1\}$ and $h = (h_\ell)_\ell$ is an element of $(\mathbb{Z}/p\mathbb{Z})^{\oplus p}$. Then we have

$$\begin{aligned} P_{c_{s,h}} &= \text{sgn}(c_{s,h}) \prod_{0 \leq \ell, m \leq p-1} \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{c_{s,h}(\ell,m) - (\ell,m) - (0,\ell k)}^{(k)}) \beta^k \right) \\ &= \text{sgn}(s) \prod_{0 \leq \ell, m \leq p-1} \left(\sum_{k=0}^{p-1} \nu_{\ell,m}(\eta_{(s(\ell) - \ell, h_\ell - \ell k)}^{(k)}) \beta^k \right) \\ &= \text{sgn}(s) \prod_{\ell=0}^{p-1} \nu_{\ell,0} \left(\prod_{m=0}^{p-1} \nu_{0,m}(Q_{s,h;\ell}) \right), \end{aligned}$$

where $Q_{s,h;\ell} = \sum_{k=0}^{p-1} \eta_{(s(\ell) - \ell, h_\ell - \ell k)}^{(k)} \beta^k$. Note that $\text{sgn}(c_{s,h}) = \text{sgn}(s)^p = \text{sgn}(s)$ since p is odd.

Since $Q_{s,h;\ell}$ is independent of m , we may prove

$$\prod_{m=0}^{p-1} \nu_{0,m}(Q_{s,h;\ell}) \equiv \varphi(Q_{s,h;\ell}) \pmod{J_2^{(n)}}$$

in the same way as Claim above.

Therefore,

$$\begin{aligned}
 & \sum_{s,h} P_{c_{s,h}} \\
 & \equiv \sum_{s,h} \operatorname{sgn}(s) \left(\prod_{\ell=0}^{p-1} \nu_{\ell,0}(\varphi(Q_{s,h;\ell})) \right) \pmod{J_2^{(n)}} \\
 & \equiv \sum_{s,h} \operatorname{sgn}(s) \left(\sum_{0 \leq k_0, \dots, k_{p-1} \leq p-1} \prod_{\ell=0}^{p-1} \varphi(\eta_{(s(\ell)-\ell, h_\ell - \ell k_\ell)}^{(k_\ell)}) \right) \pmod{J_2^{(n)}} \\
 (5.13) \quad & = \sum_{s,h,k_\ell} R_{s,h;k_0,k_1,\dots,k_{p-1}}
 \end{aligned}$$

where $R_{s,h;k_0,\dots,k_{p-1}} = \operatorname{sgn}(s) \prod_{\ell=0}^{p-1} \varphi(\eta_{(s(\ell)-\ell, h_\ell - \ell k_\ell)}^{(k_\ell)})$. We use the fact that $\nu_{\ell,0} \circ \varphi = \varphi$ (recall $\varphi(\mathbb{Z}_p[U_2^{(n)}/V_2]) \subseteq \mathbb{Z}_p[\Gamma/\Gamma^{(n)}]$) for the second congruence.

If $s(\ell) - \ell = \lambda$ (constant), $k_0 = k_1 = \dots = k_{p-1}$ and $h_\ell - \ell k_\ell = \mu$ (constant), that is, if $s(\ell) = s_\lambda(\ell) = \ell + \lambda$, $k_\ell = \kappa$ for each ℓ , and $h = h_{\mu,\kappa} = (\ell\kappa + \mu)_\ell$ for $0 \leq \kappa, \lambda, \mu \leq p-1$, we have $R_{s_\lambda, h_{\mu,\kappa}; \kappa, \dots, \kappa} = \varphi(\eta_{\lambda,\mu}^{(\kappa)})^p$. Note that $\operatorname{sgn}(s_\lambda) = 1$ because s_λ is a cyclic permutation of degree p .

Otherwise, set

$$\begin{aligned}
 k_\ell^{(w)} &= k_{\ell+w}, \\
 s^{(w)}(\ell) &= s(\ell+w) - w, \\
 h_\ell^{(w)} &= h_{\ell+w} - wk_{\ell+w},
 \end{aligned}$$

for each $0 \leq w \leq p-1$. Then we have

$$R_{s^{(0)}, h^{(0)}; k_0^{(0)}, \dots, k_{p-1}^{(0)}} = \dots = R_{s^{(p-1)}, h^{(p-1)}; k_0^{(p-1)}, \dots, k_{p-1}^{(p-1)}}$$

and $R_{s^{(w)}, h^{(w)}; k_0^{(w)}, \dots, k_{p-1}^{(w)}} \quad (0 \leq w \leq p-1)$ are distinct terms in the expansion (5.13). Hence we have

$$\sum_{w=0}^{p-1} R_{s^{(w)}, h^{(w)}; k_0^{(w)}, \dots, k_{p-1}^{(w)}} \in p\mathbb{Z}_p[\Gamma^{(n)}] \subseteq J_2^{(n)}.$$

Therefore we have

$$\begin{aligned}
\theta_2(\eta) &\equiv \sum_{s,h,k_w} R_{s,h;k_0,\dots,k_{p-1}} && \text{mod } J_2^{(n)} \\
&\equiv \sum_{0 \leq \lambda, \mu, \kappa \leq p-1} R_{s_\lambda, h_{\mu, \kappa}; \kappa, \kappa, \dots, \kappa} && \text{mod } J_2^{(n)} \\
&\equiv \sum_{0 \leq \lambda, \mu, \kappa \leq p-1} \varphi(\eta_{\lambda, \mu}^{(\kappa)})^p && \text{mod } J_2^{(n)} \\
&\equiv \left(\sum_{0 \leq \lambda, \mu, \kappa \leq p-1} \varphi(\eta_{\lambda, \mu}^{(\kappa)}) \beta^\kappa \alpha^\lambda \delta^\mu \right)^p && \text{mod } J_2^{(n)} \\
&\equiv \varphi \left(\sum_{0 \leq \lambda, \mu, \kappa \leq p-1} \eta_{\lambda, \mu}^{(\kappa)} \beta^\kappa \alpha^\lambda \delta^\mu \right)^p && \text{mod } J_2^{(n)} \\
&\equiv \varphi(\eta)^p && \text{mod } J_2^{(n)}.
\end{aligned}$$

Note that since $J_2^{(n)} \supseteq p\mathbb{Z}_p[U_2^{(n)}/V_2]$, we have

$$(x + y)^p \equiv x^p + y^p \pmod{J_2^{(n)}}.$$

- The congruence for θ_3 .

We may show the congruence for θ_3 by an argument similar to that for θ_2 . Moreover, the calculation for θ_3 is much simpler than the case for θ_2 since U_3 is normal in G . Therefore we omit the proof. $\square$

By Lemma 5.10 and the fact that $\varphi(\theta_0(\eta))$ is invertible, we have

$$\begin{aligned}
\frac{\theta_1(\eta)}{\varphi(\theta_0(\eta))} &\in 1 + J_1^{(n)}, & \frac{\tilde{\theta}_2(\eta)}{\varphi(\theta_0(\eta))} &\in 1 + \tilde{J}_2^{(n)}, \\
\frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p} &\in 1 + J_2^{(n)}, & \frac{\theta_3(\eta)}{\varphi(\theta_0(\eta))^{p^2}} &\in 1 + J_3^{(n)}.
\end{aligned}$$

Hence by Lemma 5.5, we have

$$\begin{aligned}
\log \frac{\theta_1(\eta)}{\varphi(\theta_0(\eta))} &\in J_1^{(n)}, & \log \frac{\tilde{\theta}_2(\eta)}{\varphi(\theta_0(\eta))} &\in \tilde{J}_2^{(n)}, \\
\log \frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p} &\in J_2^{(n)}, & \log \frac{\theta_3(\eta)}{\varphi(\theta_0(\eta))^{p^2}} &\in J_3^{(n)}.
\end{aligned}$$

But by Proposition 5.9, these elements are contained in the image of θ_i^+ , and therefore contained in $I_i^{(n)}$ (recall the definition of $I_i^{(n)}$). Moreover, since the p -adic logarithmic homomorphisms induce isomorphisms $1 + I_i^{(n)} \xrightarrow{\cong} I_i^{(n)}$ (by Lemma 5.6), we may conclude that

$$\begin{aligned}
(5.14) \quad & \frac{\theta_1(\eta)}{\varphi(\theta_0(\eta))} \in 1 + I_1^{(n)}, & \frac{\tilde{\theta}_2(\eta)}{\varphi(\theta_0(\eta))} &\in 1 + \tilde{I}_2^{(n)} \\
& \frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p} \in 1 + I_2^{(n)}, & \frac{\theta_3(\eta)}{\varphi(\theta_0(\eta))^{p^2}} &\in 1 + I_3^{(n)}.
\end{aligned}$$

Since each θ_i is a norm map in K -theory, it is clear that $(\theta_i(\eta))_i$ satisfies norm relations. This and (5.14) imply that $\theta(\eta) \in \Psi^{(n)}$.

5.3. Surjectivity for finite quotients. Let $(\eta_i)_i$ be an arbitrary element of $\Psi^{(n)}$. By the congruences in Definition 5.3 and by the fact that $\varphi(\eta_0)$ is invertible, we have

$$\begin{aligned} \frac{\eta_1}{\varphi(\eta_0)} &\in 1 + I_1^{(n)}, & \frac{\tilde{\eta}_2}{\varphi(\eta_0)} &\in 1 + \tilde{I}_2^{(n)}, \\ \frac{\eta_2}{\varphi(\eta_0)^p} &\in 1 + I_2^{(n)}, & \frac{\eta_3}{\varphi(\eta_0)^{p^2}} &\in 1 + I_3^{(n)}. \end{aligned}$$

Then we obtain

$$\begin{aligned} \log \frac{\eta_1}{\varphi(\eta_0)} &\in I_1^{(n)}, & \log \frac{\tilde{\eta}_2}{\varphi(\eta_0)} &\in \tilde{I}_2^{(n)}, \\ \log \frac{\eta_2}{\varphi(\eta_0)^p} &\in I_2^{(n)}, & \log \frac{\eta_3}{\varphi(\eta_0)^{p^2}} &\in I_3^{(n)} \end{aligned}$$

via the p -adic logarithms (Lemma 5.6).

Note that since $\varphi(\eta_0)$ is contained in the center of $\mathbb{Z}_p[U_i^{(n)}/V_i]^\times$, it is easy to see that

$$\begin{aligned} \text{Nr}_{\mathbb{Z}_p[\widetilde{U_2^{(n)}}/\widetilde{V_2}]/\mathbb{Z}_p[U_2^{(n)}/\widetilde{V_2}]} \varphi(\eta_0) &= \varphi(\eta_0)^p, \\ \text{Nr}_{\mathbb{Z}_p[U_1^{(n)}/\widetilde{V_2}]/\mathbb{Z}_p[U_1^{(n)} \cap \widetilde{U_2^{(n)}}/\widetilde{V_2}]} \varphi(\eta_0) &= \text{Nr}_{\mathbb{Z}_p[\widetilde{U_2^{(n)}}/\widetilde{V_2}]/\mathbb{Z}_p[U_1^{(n)} \cap \widetilde{U_2^{(n)}}/\widetilde{V_2}]} \varphi(\eta_0) \\ &= \varphi(\eta_0)^p, \\ \text{Nr}_{\mathbb{Z}_p[U_1^{(n)}/V_1]/\mathbb{Z}_p[U_3^{(n)}/V_1]} \varphi(\eta_0) &= \varphi(\eta_0)^{p^2}, \\ \text{Nr}_{\mathbb{Z}_p[U_2^{(n)}/V_2]/\mathbb{Z}_p[U_3^{(n)}/V_2]} \varphi(\eta_0)^p &= \varphi(\eta_0)^{p^2}. \end{aligned}$$

By using Lemma 5.8,

$$\begin{aligned} (y_1, \tilde{y}_2, y_2, y_3) &= \left(\log \frac{\eta_1}{\varphi(\eta_0)}, \log \frac{\tilde{\eta}_2}{\varphi(\eta_0)}, \log \frac{\eta_2}{\varphi(\eta_0)^p}, \log \frac{\eta_3}{\varphi(\eta_0)^{p^2}} \right) \\ &\in I_1^{(n)} \times \tilde{I}_2^{(n)} \times I_2^{(n)} \times I_3^{(n)} \end{aligned}$$

satisfies the trace relations in Definition 4.2 (except for $\mathbb{Z}_p[U_0^{(n)}/V_0]$ -part).

Now set $y_0 = \Gamma_{U_0^{(n)}/V_0}(\eta_0) = (1/p) \log(\eta_0^p/\varphi(\eta_0)) \in \mathbb{Z}_p[U_0^{(n)}/V_0]$. Then we may easily check that $(y_0, y_1, \tilde{y}_2, y_2, y_3)$ satisfies the trace relations in Definition 4.2. Hence we have $(y_0, y_1, \tilde{y}_2, y_2, y_3) \in \Omega^{(n)}$.

By the additive theta isomorphism (Proposition-Definition 4.3), there exists a unique element $y \in \mathbb{Z}_p[\text{Conj}(G^{(n)})]$ satisfying

$$\begin{aligned} (5.15) \quad \theta^+(y) &= (y_0, y_1, \tilde{y}_2, y_2, y_3) \\ &= \left(\Gamma_{U_0^{(n)}/V_0}(\eta_0), \log \frac{\eta_1}{\varphi(\eta_0)}, \log \frac{\tilde{\eta}_2}{\varphi(\eta_0)}, \log \frac{\eta_2}{\varphi(\eta_0)^p}, \log \frac{\eta_3}{\varphi(\eta_0)^{p^2}} \right). \end{aligned}$$

Proposition 5.11. $\omega_{G^{(n)}}(y) = 1$, that is, y is in the image of the integral logarithmic homomorphism $\Gamma_{G^{(n)}}$ where $\omega_{G^{(n)}}$ is defined as in Theorem 1.30.

Proof. By the definition, $\omega_{G^{(n)}}$ is decomposed as follows:

$$\begin{array}{ccc} \mathbb{Z}_p[\text{Conj}(G^{(n)})] & \xrightarrow{\omega_{G^{(n)}}} & G^{\text{ab}} = U_0^{(n)}/V_0 \\ \pi^{(n)} \downarrow & \nearrow \omega_{U_0^{(n)}/V_0} & \\ \mathbb{Z}_p[U_0^{(n)}/V_0] & & \end{array}$$

where $\pi^{(n)}$ is the canonical surjection. Note that $\pi^{(n)}$ is the same map as θ_0^+ by the definition of θ^+ . Therefore, we have

$$\begin{aligned} \omega_{G^{(n)}}(y) &= \omega_{U_0^{(n)}/V_0}(\theta_0^+(y)) \\ &= \omega_{U_0^{(n)}/V_0}(\Gamma_{U_0^{(n)}/V_0}(\eta_0)) \quad \text{by (5.15)} \\ &= 1 \quad \text{by Theorem 1.30 .} \end{aligned}$$

□

Let η be an element of $K_1(\mathbb{Z}_p[G^{(n)}])$ satisfying $\Gamma_{G^{(n)}}(\eta) = y$. Then η is determined up to multiplication by a torsion element of $K_1(\mathbb{Z}_p[G^{(n)}])$ (See Theorem 1.30). By the definition of η and the equation (5.15),

$$\theta^+ \circ \Gamma_{G^{(n)}}(\eta) = (y_0, y_1, \tilde{y}_2, y_2, y_3).$$

Combining with Proposition 5.9, we have

$$y_0 = \frac{1}{p} \log \frac{\eta_0^p}{\varphi(\eta_0)} = \frac{1}{p} \log \frac{\theta_0(\eta)^p}{\varphi(\theta_0(\eta))},$$

that is,

$$(5.16) \quad \Gamma_{U_0^{(n)}/V_0}(\eta_0) = \Gamma_{U_0^{(n)}/V_0}(\theta_0(\eta)).$$

We also have

$$(5.17) \quad \begin{aligned} \log \frac{\eta_1}{\varphi(\eta_0)} &= \log \frac{\theta_1(\eta)}{\varphi(\theta_0(\eta))}, & \log \frac{\tilde{\eta}_2}{\varphi(\eta_0)} &= \log \frac{\tilde{\theta}_2(\eta)}{\varphi(\theta_0(\eta))}, \\ \log \frac{\eta_2}{\varphi(\eta_0)^p} &= \log \frac{\theta_2(\eta)}{\varphi(\theta_0(\eta))^p}, & \log \frac{\eta_3}{\varphi(\eta_0)^{p^2}} &= \log \frac{\theta_3(\eta)}{\varphi(\theta_0(\eta))^{p^2}}. \end{aligned}$$

Proposition 5.12. *There exists an element $\tau \in \mu_{p-1}(\mathbb{Z}_p) \times (G^{(n)})^{\text{ab}}$ such that*

$$\theta(\eta\tau) = (\eta_0, \eta_1, \tilde{\eta}_2, \eta_2, \eta_3)$$

where $\mu_{p-1}(\mathbb{Z}_p)$ is the multiplicative group generated by all $(p-1)$ -th roots of 1 in $\mathbb{Z}_p$.

In particular, θ surjects on $\Psi^{(n)}$.

Proof. From the equation (5.16) and the fact that the integral logarithm homomorphism is injective modulo torsion elements, there exists an element $\tau \in K_1(\mathbb{Z}_p[U_0^{(n)}/V_0])_{\text{tors}}$ satisfying $\eta_0 = \tau\theta_0(\eta)$.

Then by Lemma 1.33 (3), τ is contained in

$$K_1(\mathbb{Z}_p[U_0^{(n)}/V_0])_{\text{tors}} = \mu_{p-1}(\mathbb{Z}_p) \times (U_0^{(n)}/V_0) = \mu_{p-1}(\mathbb{Z}_p) \times (G^{(n)})^{\text{ab}}.$$

Hence we have

$$\theta_0(\eta\tau) = \theta_0(\eta)\theta_0(\tau) = \theta_0(\eta)\tau = \eta_0.$$

Since $\tau \in \text{Ker}(\Gamma_{G^{(n)}})$, we have $\theta^+ \circ \Gamma_{G^{(n)}}(\eta\tau) = \theta^+ \circ \Gamma_{G^{(n)}}(\eta)$. Therefore we may replace η by $\eta\tau$ in the right hand sides of the equations (5.17).

From this fact and Lemma 5.6, we have $\theta_i(\eta\tau) = \eta_i$ for each i , using $\theta_0(\eta\tau) = \eta_0$. $\square$

5.4. Taking the projective limit. In this subsection, we take the projective limit of the surjections

$$\theta^{(n)}: K_1(\mathbb{Z}_p[G^{(n)}]) \longrightarrow \Psi^{(n)}$$

obtained by the argument in the previous subsections, and construct the surjection (5.2).

First, we study the kernel of the surjection $\theta^{(n)}$ precisely.

Suppose that $\tau \in \text{Ker}(\theta^{(n)})$. Then we have $\theta^{+, (n)} \circ \Gamma_{G^{(n)}}(\tau) = 0$ by Proposition 5.9. Since $\theta^{+, (n)}$ is an isomorphism by Proposition-Definition 4.3, we obtain $\Gamma_{G^{(n)}}(\tau) = 0$. In other words,

$$\begin{aligned} \text{Ker}(\theta^{(n)}) &\subseteq \text{Ker}(\Gamma_{G^{(n)}}) = K_1(\mathbb{Z}_p[G^{(n)}])_{\text{tors}} \\ &= \mu_{p-1}(\mathbb{Z}_p) \times (G^{(n)})^{\text{ab}} \times SK_1(\mathbb{Z}_p[G^{(n)}]). \end{aligned}$$

By the definition of SK_1 groups, norm maps of K_1 -groups induce homomorphisms on SK_1 , so we have

$$\theta^{(n)}(SK_1(\mathbb{Z}_p[G^{(n)}])) \subseteq \prod_i SK_1(\mathbb{Z}_p[U_i^{(n)}/V_i]).$$

However, $SK_1(\mathbb{Z}_p[U_i^{(n)}/V_i]) = 1$ since each $U_i^{(n)}/V_i$ is abelian (Proposition 1.33 (3)). Hence we obtain $SK_1(\mathbb{Z}_p[G^{(n)}]) \subseteq \text{Ker}(\theta^{(n)})$.

On the other hand, for every element $\tau \in \mu_{p-1}(\mathbb{Z}_p) \times (G^{(n)})^{\text{ab}}$ not equal to 1, we have $\theta_0^{(n)}(\tau) = \tau \neq 1$. This implies $(\mu_{p-1}(\mathbb{Z}_p) \times (G^{(n)})^{\text{ab}}) \cap \text{Ker}(\theta^{(n)}) = \{1\}$.

Therefore we may conclude that $\text{Ker}(\theta^{(n)}) = SK_1(\mathbb{Z}_p[G^{(n)}])$, and we obtain an exact sequence of projective systems of abelian groups

$$1 \longrightarrow SK_1(\mathbb{Z}_p[G^{(n)}]) \longrightarrow K_1(\mathbb{Z}_p[G^{(n)}]) \xrightarrow{\theta^{(n)}} \Psi^{(n)} \longrightarrow 1.$$

By Proposition 1.33 (2), $SK_1(\mathbb{Z}_p[G^{(n)}])$ is a finite abelian group for every $n \geq 1$. Hence the projective system $\{SK_1(\mathbb{Z}_p[G^{(n)}])\}_{n \in \mathbb{N}}$ satisfies the Mittag-Leffler condition, which implies $\varprojlim_n SK_1(\mathbb{Z}_p[G^{(n)}]) = 0$. By taking the projective limit of (5.18), we obtain the following exact sequence

$$(5.19) \quad 1 \rightarrow \varprojlim_n SK_1(\mathbb{Z}_p[G^{(n)}]) \rightarrow \varprojlim_n K_1(\mathbb{Z}_p[G^{(n)}]) \xrightarrow{(\theta^{(n)})_n} \Psi \rightarrow 1.$$

Therefore, the surjectivity of (5.2) is reduced to the following proposition.

Proposition 5.13. *The homomorphism*

$$K_1(\Lambda(G)) \rightarrow \varprojlim_n K_1(\mathbb{Z}_p[G^{(n)}])$$

induced by canonical homomorphisms $K_1(\Lambda(G)) \rightarrow K_1(\mathbb{Z}_p[G^{(n)}])$ is an isomorphism.

Proof. Apply Proposition 1.8 for $\Lambda(G)$ and $\{\mathbb{Z}_p[G^{(n)}]\}_{n \in \mathbb{N}}$. $\square$

Remark 5.14. In the case of Kato's Heisenberg type ([Kato1]), that is, in the case that $\text{Gal}(F^\infty/F) \cong \overline{G}$ where

$$\overline{G} = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix} \times \Gamma = \overline{G}^f \times \Gamma,$$

it is known ([Oliver]) that

$$SK_1(\mathbb{Z}_p[\overline{G}^{(n)}]) = SK_1(\mathbb{Z}_p[\overline{G}^f]) \oplus SK_1(\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]) = 1,$$

therefore we may show that the theta map

$$K_1(\Lambda(\overline{G})) \xrightarrow{\cong} \overline{\Psi} \left(\subseteq \prod_i \Lambda(U_i/V_i) \right)$$

is an isomorphism, and $\xi_{F^\infty/F}$ is determined uniquely.

Remark 5.15. By the calculation above, we know that the kernel of the theta map tends to be $SK_1(\Lambda(G))$. It is conjectured that $SK_1(\Lambda(G))$ vanishes for general compact p -adic Lie groups.

For our special case $G = G^f \times \Gamma$, Otmar Venjakob announced to the author that he and Peter Schneider have recently proven the vanishing of $SK_1(\Lambda(G))$. If we admit their result, we may prove the uniqueness of the p -adic zeta function $\xi_{F^\infty/F}$ which we would construct in §8.

6. LOCALIZED THETA MAP

In this section, we construct *the localized theta map*

$$\theta_S : K_1(\Lambda(G)_S) \longrightarrow \Psi_S.$$

Let $\Lambda(\Gamma) = \mathbb{Z}_p[[\Gamma]]$ be the Iwasawa algebra for Γ . Then since $G = G^f \times \Gamma$, we have

$$\begin{aligned} \Lambda(G) &= \mathbb{Z}_p[[G^f \times \Gamma]] \\ &\cong \mathbb{Z}_p[G^f] \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[\Gamma]] \\ &\cong \Lambda(\Gamma)[G^f], \end{aligned}$$

and we also have $\Lambda(U_i/V_i) \cong \Lambda(\Gamma)[U_i^f/V_i^f]$

Let S be the canonical Ore set (see §2.1) for the group G and let S_0 be that for Γ . Note that $S_0 = \Lambda(\Gamma) \setminus p\Lambda(\Gamma)$ (see Example 2.7).

Lemma 6.1. $\Lambda(G)_S \cong \Lambda(\Gamma)_{S_0}[G^f]$ and $\Lambda(U_i/V_i) \cong \Lambda(\Gamma)_{S_0}[U_i^f/V_i^f]$.

Proof. We show this lemma only for the group G .

First, we show that $\Lambda(G)_S = \Lambda(G)_{S_0}$ (by abuse of notation, we also denote the image of S_0 under the canonical map $\Lambda(\Gamma) \longrightarrow \Lambda(G)$ by the same symbol S_0). For an arbitrary $f_0 \in S_0$, $\Lambda(G)/\Lambda(G)f_0 \cong (\Lambda(\Gamma)/\Lambda(\Gamma)f_0)[G^f]$ is $\mathbb{Z}_p$ -finitely generated, so $S_0 \subseteq S$. Then by the universality of Ore localizations (Proposition-Definition 1.13), we obtain the canonical homomorphism

$$(6.1) \quad \Lambda(G)_{S_0} \longrightarrow \Lambda(G)_S.$$

Let f be an element of S . By the definition of the canonical Ore set, $\Lambda(G)/\Lambda(G)f$ is a finitely generated left $\mathbb{Z}_p$ -module.⁷ Let $\mathfrak{z}$ be a kernel of the following composition of homomorphisms:

$$(6.2) \quad \Lambda(\Gamma) \longrightarrow \Lambda(G) \longrightarrow \Lambda(G)/\Lambda(G)f,$$

where the first map is the canonical map

$$\Lambda(\Gamma) \longrightarrow \Lambda(G) \cong \Lambda(\Gamma) \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[G^f]; f_0 \mapsto f_0 \otimes 1.$$

Take an arbitrary non-zero element $g_0 \in \mathfrak{z}$. Since $\Lambda(\Gamma)$ is a unique factorization domain, g_0 has a decomposition

$$g_0 = up^n \pi_1^{n_1} \cdots \pi_k^{n_k},$$

where $u \in \Lambda(\Gamma)^\times$ and π_i is a prime element of $\Lambda(\Gamma)$.

Suppose that for every non-zero $g_0 \in \mathfrak{z}$, the index of the prime p (n in the decomposition above) is not zero. Then we have $\mathfrak{z} \subseteq p\Lambda(\Gamma)$, and this induces a surjection $\Lambda(\Gamma)/\mathfrak{z} \rightarrow \Lambda(\Gamma)/p\Lambda(\Gamma)$. This is contradiction since $\Lambda(\Gamma)/\mathfrak{z} \subseteq \Lambda(G)/\Lambda(G)f$ is finitely generated over $\mathbb{Z}_p$ by the definition of S , but $\Lambda(\Gamma)/p\Lambda(\Gamma) \cong \mathbb{F}_p[[\Gamma]]$ is not so. Hence there exists $g_0 \in \mathfrak{z}$ such that $p \nmid g_0$, which implies $g_0 \in S_0$.

Via the homomorphism (6.2), g_0 is contained in $\Lambda(G)f$, hence there exists $\phi \in \Lambda(G)$ satisfying $g_0 = \phi f$. This means $(g_0^{-1}\phi) \cdot f = 1$ in $\Lambda(G)_{S_0}$, in other words, f has its (left) inverse element $g_0^{-1}\phi$ in $\Lambda(G)_{S_0}$. Hence, by the universality of Ore localizations again, we obtain

$$(6.3) \quad \Lambda(G)_S \longrightarrow \Lambda(G)_{S_0}.$$

It is easy to show that the homomorphisms (6.1) and (6.3) are the inverse maps of each other, therefore $\Lambda(G)_S \cong \Lambda(G)_{S_0}$.

On the other hand, the canonical homomorphisms

$$\Lambda(G)_{S_0} \longrightarrow \Lambda(\Gamma)_{S_0}[G^f]$$

obtained by the universality of Ore localizations and

$$\Lambda(\Gamma)_{S_0}[G^f] = \Lambda(\Gamma)_{S_0} \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[G^f] \longrightarrow \Lambda(G)_{S_0}$$

obtained by the universality of tensor products from the map

$$\Lambda(\Gamma)_{S_0} \times \mathbb{Z}_p[G^f] \rightarrow \Lambda(G)_{S_0}; (\phi_0, h) \mapsto \phi_0 h$$

are inverse maps of each other. This implies $\Lambda(G)_S \cong \Lambda(\Gamma)_{S_0}[G^f]$.

The result for each $\Lambda(U_i/V_i)_S$ is obtained in the same way.

□

By this lemma, we may replace the coefficient ring $\mathbb{Z}_p$ by $\Lambda(\Gamma)_{S_0}$ and apply the same discussion in §4 and §5.

Since we need to use p -adic logarithms to translate the additive theta map, we take the p -adic completion $\widehat{\Lambda(\Gamma)_{S_0}}$ of $\Lambda(\Gamma)_{S_0}$. Set $R = \widehat{\Lambda(\Gamma)_{S_0}}$. Then the additive theta map

$$\widehat{\theta}^+ : R[\text{Conj}(G^f)] \xrightarrow{\sim} \widehat{\Omega}$$

⁷Since H is finite, a left $\mathbb{Z}_p[H]$ -module is finitely generated if and only if it is finitely generated as a left $\mathbb{Z}_p$ -module.

is obtained by the completely same argument as in §4, where $\widehat{\Omega}$ is the R -submodule of $\prod_i R[U_i^f/V_i^f]$ defined by the same conditions as in Definition

4.2. More precisely, $\widehat{\Omega}$ is an R -submodule consisting of all elements $(\widehat{y}_i)_i$ satisfying the trace relations and $\widehat{y}_i \in \widehat{I}_i$ for each i where $\widehat{I}_i = I_i^f \otimes_{\mathbb{Z}_p} R$.

We use the generalized integral p -adic logarithm (See §1.4)

$$\Gamma_{R[G^f], J}: K_1(R[G^f], J) \longrightarrow R[\text{Conj}(G^f)],$$

where J is the Jacobson radical of $R[G^f]$. In this case, J is the kernel of the natural surjection $R[G^f] \rightarrow R/pR$. By the same argument as in §5, we have

$$\widehat{\theta}^+ \circ \Gamma_{R[G^f], J}(\widehat{\eta}') \in \widehat{\Omega},$$

for an arbitrary $\widehat{\eta}' \in K_1(R[G^f], J)$, and we may conclude that $\widehat{\theta}(\widehat{\eta}')$ is contained in $\widehat{\Psi}$, where $\widehat{\Psi}$ is the subgroup of $\prod_i R[U_i^f/V_i^f]$ consisting of all elements $(\widehat{\eta}'_i)_i$ which satisfy the norm relations and the following congruences:

$$(6.4) \quad \begin{aligned} \widehat{\eta}'_1 &\equiv \varphi(\widehat{\eta}'_0) \pmod{\widehat{I}_1}, & \widehat{\eta}'_2 &\equiv \varphi(\widehat{\eta}'_0) \pmod{\widehat{I}_2}, \\ \widehat{\eta}'_2 &\equiv \varphi(\widehat{\eta}'_0)^p \pmod{\widehat{I}_2}, & \widehat{\eta}'_3 &\equiv \varphi(\widehat{\eta}'_0)^{p^2} \pmod{\widehat{I}_3}. \end{aligned}$$

Let $\widehat{\eta}$ be an arbitrary element of $K_1(R[G^f])$. We use the same notation $\widehat{\eta}$ for its lift to $R[G^f]^\times$. Let $F \subseteq R[G^f]$ be a subset of representatives of $\pi^\times: R[G^f]^\times \rightarrow (R[G^f]/J)^\times \cong \mathbb{F}_p((T))^\times$, that is, F be a subset of $R[G^f]$ for which $\pi^\times$ induces a bijection of sets $F \xrightarrow{\sim} (R[G^f]/J)^\times$. Note that we can take F as a subset of $R^\times$.

Then there exists $\phi \in F$ and $\widehat{\eta}' \in 1 + J$ such that $\widehat{\eta} = \phi \widehat{\eta}'$. $\widehat{\eta}'$ defines an element of $K_1(R[G^f], J)$, and we have already known that $\widehat{\theta}(\widehat{\eta}')$ is contained in $\widehat{\Psi}$. Therefore, if we prove that $\widehat{\theta}(\phi) \in \widehat{\Psi}$ for every $\phi \in F$, we may conclude that $\widehat{\theta}(K_1(R[G^f])) \subseteq \widehat{\Psi}$. It suffices to show that $\widehat{\theta}(r) \in \widehat{\Psi}$ for every $r \in R^\times$.

For an arbitrary $r \in R^\times$, we have

$$\begin{aligned} \widehat{\theta}_0(r) &= r, & \widehat{\theta}_1(r) &= \widehat{\theta}_2(r) = r^p, \\ \widehat{\theta}_2(r) &= r^{p^2}, & \widehat{\theta}_3(r) &= r^{p^3}, \end{aligned}$$

by direct calculation.⁸ This implies that $(\widehat{\theta}_i(r))_i$ satisfies the norm relations.

The congruences for $(\theta_i(r))_i$ are derived from the following relation

$$r^p \equiv \varphi(r) \pmod{pR}$$

which follows from the definition of the Frobenius automorphism (See §1.4).

Now $(\widehat{\theta}_i(r))_i \in \widehat{\Psi}$ holds.

$$\text{Set } \Psi_S = \widehat{\Psi} \cap \prod_i \Lambda(\Gamma)_{S_0}[U_i^f/V_i^f].$$

⁸Since R is in the center of $R[G^f]$, the images of norm maps of $r \in R$ are easily calculated by using the method in §1.3.

Proposition 6.2. Ψ_S is characterized as the subgroup of $\prod_i \Lambda(U_i/V_i)_S^\times$ consisting of all elements $(\eta_0, \eta_1, \tilde{\eta}_2, \eta_2, \eta_3)$ satisfying the following two conditions.

(1) (norm relations)

- (rel-1) $\text{Nr}_{\Lambda(U_0/V_0)_S/\Lambda(U_1/V_0)_S} \eta_0 \equiv \eta_1,$
 - (rel-2) $\text{Nr}_{\Lambda(U_0/V_0)_S/\Lambda(\tilde{U}_2/V_0)_S} \eta_0 \equiv \tilde{\eta}_2,$
 - (rel-3) $\text{Nr}_{\Lambda(\tilde{U}_2/\tilde{V}_2)_S/\Lambda(U_2/\tilde{V}_2)_S} \tilde{\eta}_2 \equiv \eta_2,$
 - (rel-4) $\text{Nr}_{\Lambda(U_1/\tilde{V}_2)_S/\Lambda(U_1 \cap \tilde{U}_2/\tilde{V}_2)_S} \eta_1 \equiv \text{Nr}_{\Lambda(\tilde{U}_2/\tilde{V}_2)_S/\Lambda(U_1 \cap \tilde{U}_2/\tilde{V}_2)_S} \tilde{\eta}_2,$
 - (rel-5) $\text{Nr}_{\Lambda(U_1/V_1)_S/\Lambda(U_3/V_1)_S} \eta_1 \equiv \eta_3,$
 - (rel-6) $\text{Nr}_{\Lambda(U_2/V_2)_S/\Lambda(U_3/V_2)_S} \eta_2 \equiv \eta_3.$
- (These are the same as those of Ψ . See Figure 1)

(2) (congruences)

$$\begin{aligned} \eta_1 &\equiv \varphi(\eta_0) \pmod{I_{S,1}}, & \tilde{\eta}_2 &\equiv \varphi(\eta_0) \pmod{\widetilde{I_{S,2}}}, \\ \eta_2 &\equiv \varphi(\eta_0)^p \pmod{I_{S,2}}, & \eta_3 &\equiv \varphi(\eta_0)^{p^2} \pmod{I_{S,3}}, \end{aligned}$$

where $I_{S,i} = I_i^f \otimes_{\mathbb{Z}_p} \Lambda(\Gamma)_{S_0}$.

Proof. An arbitrary element $(\eta_i)_i$ of Ψ_S satisfies the norm relations by the definition of Ψ_S . Since $(\eta_i)_i$ is contained in $\widehat{\Psi}$, it satisfies (6.4). Then the desired congruences hold (note that $\widehat{I}_i \cap \Lambda(\Gamma)_{S_0}[U_i^f/V_i^f] = I_i^f \otimes_{\mathbb{Z}_p} \Lambda(\Gamma)_{S_0} = I_{S,i}$). $\square$

Now let $\theta_S = (\theta_{S,i})_i$ be the family of the homomorphisms $\theta_{S,i}$ where each $\theta_{S,i}$ is defined as the composite of the norm map

$$\text{Nr}_{\Lambda(G)_S/\Lambda(U_i)_S} : K_1(\Lambda(G)_S) \rightarrow K_1(\Lambda(U_i)_S)$$

and the canonical homomorphism

$$K_1(\Lambda(U_i)_S) \rightarrow K_1(\Lambda(U_i/V_i)_S) = \Lambda(U_i/V_i)_S^\times$$

induced by $\Lambda(U_i)_S \rightarrow \Lambda(U_i/V_i)_S$.

Proposition 6.3. Ψ_S contains the image of θ_S .

Proof. By the commutative diagram

$$\begin{array}{ccc} K_1(\Lambda(G)_S) & \xrightarrow{\theta_S} & \prod_i \Lambda(U_i/V_i)_S^\times \\ \downarrow & & \downarrow \\ K_1(R[G^f]) & \xrightarrow{\widehat{\theta}} & \prod_i R[U_i^f/V_i^f]^\times, \end{array}$$

we have $\text{Image}(\theta_S) \subseteq \text{Image}(\widehat{\theta}) \subseteq \widehat{\Psi}$ (note that the right vertical map is injective since the canonical map $\Lambda(\Gamma)_{S_0} \rightarrow R$ is injective, using the fact that $\Lambda(\Gamma)_{S_0}$ is separated with respect to p -adic topology). Therefore,

$$\text{Image}(\theta_S) \subseteq \widehat{\Psi} \cap \left(\prod_i \Lambda(U_i/V_i)_S^\times \right) = \Psi_S.$$

$\square$

Proposition 6.4. $\Psi_S \cap \prod_i \Lambda(U_i/V_i)^\times = \Psi.$

Proof. This follows easily from the fact $I_{S,i} \cap \Lambda(U_i/V_i) = I_i.$ $\square$

We can summarize the two propositions above as follows:

Theorem 6.5. $\theta_S: K_1(\Lambda(G)_S) \longrightarrow \Psi_S$ is the localized theta map (defined in Definition 3.3) for G .

7. CONGRUENCES AMONG ABELIAN p -ADIC ZETA PSEUDOMEASURES

7.1. Norm relations. Let F_{U_i} (resp. F_{V_i}) be the maximal subfield of F^∞ fixed by U_i (resp. V_i).

In the previous sections we have constructed the theta map θ and its localized version θ_S . If the Iwasawa main conjecture is true in our case, the assumption of Theorem 3.4 should be true, that is, the family of the abelian p -adic zeta pseudomeasures $(\xi_0, \xi_1, \tilde{\xi}_2, \xi_2, \xi_3)$ (each $\xi_i \in \text{Frac}(\Lambda(U_i/V_i))$ is the p -adic zeta pseudomeasure for F_{V_i}/F_{U_i}) should be contained in Ψ_S . For $(\xi_i)_i$ to be an element of Ψ_S , it is necessary that ξ_i 's satisfy the norm relations and the congruences in Proposition 6.2.

Proposition 7.1. ξ_i 's satisfy the norm relations in Proposition 6.2.

Proof. We may show this proposition by formal calculation, using only the interpolation properties (See Definition 2.14 (2.2)). Here we only show

$$\text{Nr}_{\Lambda(U_0/V_0)_S/\Lambda(U_1/V_0)_S} \xi_0 \equiv \xi_1 \quad \text{in } \Lambda(U_1/V_0)_S.$$

Let ρ be an arbitrary Artin representation of U_1/V_0 . Then by Lemma 3.5, we have

$$\begin{aligned} (\text{Nr } \xi_0)(\rho \otimes \kappa^r) &= \xi_0(\text{Ind}_{U_1}^{U_0}(\rho \otimes \kappa^r)) \\ &= \xi_0(\text{Ind}_{U_1}^{U_0}(\rho) \otimes \kappa^r) \\ &= L_\Sigma(1-r; F_{V_0}/F_{U_0}, \text{Ind}_{U_1}^{U_0}(\rho)) \\ &= L_\Sigma(1-r; F_{V_0}/F_{U_1}, \rho) \\ &= \xi_1(\rho \otimes \kappa^r) \end{aligned}$$

for every positive integer r divisible by $p-1$.

We may also show the other norm relations by the same argument. $\square$

7.2. Kato's observation. Now let us study the congruences among ξ_i 's.

Let I_2'' be the image of the composition

$$\mathbb{Z}_p[[\text{Conj}(\widetilde{U}_2)]] \xrightarrow{\text{Tr}} \mathbb{Z}_p[[\text{Conj}(U_2)]] \rightarrow \mathbb{Z}_p[[U_2/V_2]].$$

We have the following explicit description of I_2'' :

$$I_2'' = \varprojlim_n \left([\beta^b \gamma^c h_\zeta(c \neq 0), p\beta^b \zeta^f]_{\mathbb{Z}_p[\Gamma/\Gamma^{p^n}]} \right) \subseteq \Lambda(U_2/V_2).$$

Proposition 7.2 (Congruences among abelian p -adic zeta functions). *The p -adic zeta functions ξ_i for F_{V_i}/F_{U_i} satisfy the following congruences:*

$$\begin{aligned} (1) \quad & \xi_1 \equiv \varphi(\xi_0) \pmod{I_{S,1}}, \\ (2) \quad & \widetilde{\xi_2} \equiv \varphi(\xi_0) \pmod{\widetilde{I_{S,2}}}, \\ (2) \quad & \xi_2 \equiv c_2 \pmod{I''_{S,2}}, \\ (3) \quad & \xi_3 \equiv c_3 \pmod{I_{S,3}}, \end{aligned}$$

where c_2 and c_3 are certain elements of $\Lambda(\Gamma)_{S_0}$.

Remark 7.3. These congruences are **not** sufficient to show that $(\xi_i)_i \in \Psi_S$. Hence we should modify Burns' technique (Proposition 3.4) to prove the Iwasawa main conjecture in our case. This modification is discussed in the next section.

Kato observed in [Kato1] that these kinds of congruences among abelian p -adic zeta pseudomeasures were derived from Deligne-Ribet's method of p -adic Hilbert modular forms. The fundamental philosophy of p -adic (Λ -adic) Hilbert modular forms is as follows (quite rough):

(#) Congruence between constant terms of two p -adic (Λ -adic) Hilbert modular forms is derived from congruences between coefficients of all positive degree terms of them.

Before the precise proof of Proposition 7.2, let us review how to derive the congruence between abelian p -adic zeta pseudomeasures from Hilbert modular forms, following [Kato1] §4. We take the congruence $\xi_1 \equiv \varphi(\xi_0) \pmod{I_{S,1}}$ as an example.

The $\Lambda(U_i/V_i)$ -adic F_{U_i} -Hilbert Eisenstein series

$$f_i = \frac{\xi_i}{2^{[F_{U_i}:\mathbb{Q}]}} + \sum_{(\mathfrak{a},x) \in P_i} \left(\frac{F_{V_i}/F_{U_i}}{\mathfrak{a}} \right) q^x \quad i = 0, 1$$

was essentially constructed by Deligne and Ribet, where

$$P_i = \{(\mathfrak{a}, x) \mid \mathfrak{a} \subseteq \mathcal{O}_{F_{U_i}} : \text{non-zero ideal prime to } \Sigma, \\ x \in \mathfrak{a} : \text{totally positive}\}$$

and $\left(\frac{L/K}{-} \right)$ denotes the Artin symbol in $\text{Gal}(L/K)$ for an abelian extension L/K .

By restricting f_1 to the Hilbert modular variety of F embedded as the diagonal in the Hilbert modular variety of F_{U_1} , we obtain a $\Lambda(U_1/V_1)$ -adic F -Hilbert modular form

$$(7.1) \quad g_1 = \frac{\xi_1}{2^{rp}} + \sum_{(\mathfrak{a},x) \in P_1} \left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) q^{\text{Tr}_{F_{U_1}/F}(x)}$$

where r is $[F:\mathbb{Q}]$. Note that $[F_{U_1}:\mathbb{Q}]$ is equal to rp . The Galois group

$$\text{Gal}(F_{U_1}/F) = \langle \beta \rangle$$

acts on P_1 from the left as

$$\beta^j * (\mathfrak{a}, x) = (\beta^j \mathfrak{a}, \beta^j x)$$

for each $0 \leq j \leq p-1$.

Then we may divide positive degree terms of g_1 by isotropic subgroups $\text{Gal}(F_{U_1}/F)_{(\mathfrak{a},x)}$ as follows:

$$g_1 = \frac{\xi_1}{2^{rp}} + \sum_{(\mathfrak{a},x) \in P'_1} \left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) q^{\text{Tr}_{F_{U_1}/F}(x)} + \sum_{(\mathfrak{a},x) \in P''_1} \left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) q^{\text{Tr}_{F_{U_1}/F}(x)}$$

where

$$P'_1 = \{(\mathfrak{a}, x) \in P_1 \mid \text{Gal}(F_{U_1}/F)_{(\mathfrak{a},x)} = \text{Gal}(F_{U_1}/F)\},$$

$$P''_1 = \{(\mathfrak{a}, x) \in P_1 \mid \text{Gal}(F_{U_1}/F)_{(\mathfrak{a},x)} = \{\text{id}_{F_{U_1}}\}\}.$$

One observes that there exists a bijection $P_0 \xrightarrow{\sim} P'_1; (\mathfrak{b}, y) \mapsto (\mathfrak{a}, x)$ where $\mathfrak{a} = \mathfrak{b}\mathcal{O}_{F_{U_1}}$ and $x = y$. By the functoriality of the Artin symbol, we have

$$g_1 = \frac{\xi_1}{2^{rp}} + \sum_{(\mathfrak{b},y) \in P_0} \text{Ver} \left(\left(\frac{F_{V_0}/F_{U_0}}{\mathfrak{b}} \right) \right) q^{py} + \sum_{(\mathfrak{a},x) \in P''_1} \left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) q^{\text{Tr}_{F_{U_1}/F}(x)},$$

where $\text{Ver}: \text{Gal}(F_{V_0}/F_{U_0}) \rightarrow \text{Gal}(F_{V_1}/F_{U_1})$ is the Verlagerung (transfer) homomorphism.

Lemma 7.4. *Let H_1^f and H_2^f be arbitrary subgroups of G^f . Set $H_i = H_i^f \rtimes \Gamma$. If H_1 contains H_2 , the Verlagerung homomorphism*

$$\text{Ver}_{H_2}^{H_1}: H_1^{\text{ab}} \rightarrow H_2^{\text{ab}}$$

coincides with the composition of the Frobenius homomorphism $\varphi: H_1^{\text{ab}} \rightarrow \Gamma$ and the canonical injection $\Gamma \rightarrow H_2^{\text{ab}}$.

Proof. By the transitivity of the Verlagerung homomorphism, it suffices to prove the proposition when $(H_1 : H_2) = p$. In this case it is well known that H_2^f is normal in H_1^f . Let σ be an element of H_1^f which generates H_1^f/H_2^f . Then $\{\sigma^j\}_{j=0}^{p-1}$ is a set of representatives of H_1/H_2 . Suppose that $h \in H_1^f$ satisfies $h \equiv \sigma^{j(h)} \pmod{H_2^f}$. By the definition of the Verlagerung homomorphisms,

$$\text{Ver}_{H_2}^{H_1}(h) = \prod_{j=0}^{p-1} \sigma^{-(j+j(h))} h \sigma^j = h^p = 1.$$

□

Hence in our special case (the case when the exponent of G^f is p), we may identify every Verlagerung homomorphism with the Frobenius homomorphism φ .

If we set $\varphi(q^y) = q^{py}$, we have

$$g_1 = \frac{\xi_1}{2^{rp}} + \sum_{(b,y) \in P_0} \varphi \left(\left(\frac{F_{V_0}/F}{b} \right) q^y \right) + \sum_{(a,x) \in P_1''} \left(\frac{F_{V_1}/F_{U_1}}{a} \right) q^{\text{Tr}_{F_{U_1}/F}(x)}$$

and the first summation is the same as positive degree terms of $\varphi(f_0)$.

Next, let $\tilde{P}_1''$ be a system of representatives of the orbital decomposition $\text{Gal}(F_{U_1}/F_{U_0}) \backslash P_1''$. By the property of the Artin symbol, we have

$$\sum_{j=0}^{p-1} \left(\frac{F_{V_1}/F_{U_1}}{\beta^j c} \right) = \sum_{j=0}^{p-1} \beta^j \left(\frac{F_{V_1}/F_{U_1}}{c} \right) \beta^{-j},$$

therefore the right hand side of the equation above is contained in I_1 since I_1 is the image of θ_1^+ (trace map).

Hence, we obtain

$$g_1 - \varphi(f_0) = \left(\frac{\xi_1}{2^{rp}} - \frac{\varphi(\xi_0)}{2^r} \right) + \sum_{(c,z) \in \tilde{P}_1''} \left\{ \sum_{j=0}^{p-1} \beta^j \left(\frac{F_{V_1}/F_{U_1}}{c} \right) \beta^{-j} \right\} q^{\text{Tr}_{F_{U_1}/F}(z)}$$

and all positive degree terms of $g_1 - \varphi(f_0)$ are contained in I_1 , which implies

$$\frac{\xi_1}{2^{rp}} \equiv \frac{\varphi(\xi_0)}{2^r} \pmod{I_{S,1}}$$

by the philosophy $(\sharp)$.

Note that $2^p \equiv 2 \pmod{I_{S,1}}$ since $p \in I_{S,1}$. Hence we have the desired congruence.

In the rest of this section, we make this argument precise and prove the congruences in Proposition 7.2. We adopt the method of Ritter and Weiss ([R-W3]).

7.3. Approximation of abelian p -adic zeta pseudomeasures. First, following [R-W3], we approximate the abelian p -adic zeta pseudomeasures and reduce the congruences among pseudomeasures to those among special values of partial zeta functions.

Let W_i be U_i/V_i , and let W_i^f be U_i^f/V_i . For an arbitrary open set $\mathcal{U} \subseteq W_i$, we define a non-negative integer $m(\mathcal{U})$ by $\kappa^{p-1}(\mathcal{U}) = 1 + p^{m(\mathcal{U})} \mathbb{Z}_p$ where κ is the p -adic cyclotomic character. Let $\omega = m(\Gamma)$. Then one can easily see that $m(\mathcal{U}^f \times \Gamma^{p^j}) = \omega + j$ for an arbitrary $\mathcal{U}^f \subseteq W_i^f$ and $j \in \mathbb{Z}_{\geq 0}$.

Lemma 7.5. *The canonical surjection*

$$\mathbb{Z}_p[[W_i]] \rightarrow \mathbb{Z}_p[W_i/\mathcal{U}]/p^{m(\mathcal{U})} \mathbb{Z}_p[W_i/\mathcal{U}]$$

induces the isomorphism

$$\mathbb{Z}_p[[W_i]] \xrightarrow{\sim} \varprojlim_{\mathcal{U} \subseteq W_i: \text{ open}} \mathbb{Z}_p[W_i/\mathcal{U}]/p^{m(\mathcal{U})} \mathbb{Z}_p[W_i/\mathcal{U}].$$

Proof. The injectivity is obvious since $m(\mathcal{U})$ is not bounded. The surjectivity is also not difficult, for we can construct the lift of $(x_{\mathcal{U}})_{\mathcal{U}}$ to $\mathbb{Z}_p[[W_i]]$. See [Kakde] §4.2.1. $\square$

It is clear that $\{\Gamma^{p_j}\}_{j \in \mathbb{Z}_{\geq 0}}$ is the cofinal system of the inverse limit above.

Let $\varepsilon = \varepsilon_i$ is a $\mathbb{C}$ -valued locally constant function on W_i . Then there exists an open subset $\mathcal{U} \subseteq W_i$ such that ε is constant on each coset of $W_i/\mathcal{U}$. Therefore we can write

$$\varepsilon = \sum_{x \in W_i/\mathcal{U}} \varepsilon(x) \delta^{(x)}$$

where $\delta^{(x)}$ is the characteristic function with respect to a coset x , that is,

$$\delta^{(x)}(w) = \begin{cases} 1 & \text{if } w \in x, \\ 0 & \text{otherwise.} \end{cases}$$

Definition 7.6 (partial zeta function). Let x be an arbitrary coset in $W_i/\mathcal{U}$. Then we call

$$\zeta_{F_{V_i}/F_{U_i}}(s, \delta^{(x)}) = \sum_{0 \neq \mathfrak{a} \subseteq \mathcal{O}_{F_{U_i}}} \frac{\delta^{(x)}\left(\left(\frac{F_{V_i}/F_{U_i}}{\mathfrak{a}}\right)\right)}{(N\mathfrak{a})^s}$$

the partial zeta function for $\delta^{(x)}$. Here $\left(\frac{F_{V_i}/F_{U_i}}{-}\right)$ is the Artin symbol and $N\mathfrak{a}$ is the absolute norm of an ideal $\mathfrak{a}$. This function has analytic continuation to the whole complex plane except for a simple pole at $s = 1$, and for every $k \in \mathbb{N}$, $\zeta_{F_{V_i}/F_{U_i}}(1 - k, \delta^{(x)})$ is a rational number.

For an arbitrary local constant function ε on W_i , we define the partial zeta function for ε as

$$\zeta_{F_{V_i}/F_{U_i}}(s, \varepsilon) = \sum_{x \in W_i/\mathcal{U}} \varepsilon(x) \zeta_{F_{V_i}/F_{U_i}}(s, \delta^{(x)})$$

where $\varepsilon = \sum_{x \in W_i/\mathcal{U}} \varepsilon(x) \delta^{(x)}$ is the decomposition as above.

For an arbitrary element $w \in W_i$, $\mathbb{Q}_p$ -valued locally constant function ε on W_i , and positive integer k divisible by $p - 1$, we define

$$\Delta_i^w(1 - k, \varepsilon) = \zeta_{F_{V_i}/F_{U_i}}(1 - k, \varepsilon) - \kappa(w)^k \zeta_{F_{V_i}/F_{U_i}}(1 - k, \varepsilon_w) \in \mathbb{Q}_p$$

where $\varepsilon_w(w') = \varepsilon(ww')$ for every $w' \in W_i$. Deligne and Ribet showed the integrality of $\Delta_i^w(1 - k, \delta^{(x)})$, that is, for an arbitrary $w \in W_i$ and an arbitrary coset $x \in W_i/\mathcal{U}$,

$$\Delta_i^w(1 - k, \delta^{(x)}) \in \mathbb{Z}_p$$

([De-Ri] Theoreme (0.4), see also Hypothesis (H_{n-1}) in [Coates]).

Proposition 7.7 (Approximation lemma, Ritter-Weiss). *Let $\mathcal{U} \subseteq W_i$ be an arbitrary open set. Then for every positive integer k divisible by $p - 1$ and $w \in W_i$, $(1 - w)\xi_i$ maps to*

$$(7.2) \quad \sum_{x \in W_i/\mathcal{U}} \Delta_i^w(1 - k, \delta^{(x)}) \kappa(x)^{-k} x \quad \text{mod } p^{m(\mathcal{U})}$$

under the canonical surjection $\mathbb{Z}_p[[W_i]] \rightarrow \mathbb{Z}_p[W_i/\mathcal{U}]/p^{m(\mathcal{U})}\mathbb{Z}_p[W_i/\mathcal{U}]$. (Note that $\kappa(x)^{-k}$ is well-defined by the definition of $m(\mathcal{U})$.)

Sketch of the proof. Let ξ_i^w be the limit element of (7.2). This limit element is independent of k due to Deligne-Ribet's congruence ([De-Ri] Theoreme (0.4) and [Coates] Hypothesis (C_0)). One sees that ξ_i^w is contained in $\mathbb{Z}_p[[W_i]]$ (that is, ξ_i^w is a p -adic measure on W_i) by the integrality of $\Delta_i^w(1-k, \delta^{(x)})$ and Lemma 7.5.

Then one sees that

$$\int_{W_i} \varepsilon \kappa^k d\xi_i^w = \Delta_i^w(1-k, \varepsilon)$$

for each locally constant function ε on W_i and positive integer k divisible by $p-1$, but this property characterizes the p -adic measure $(1-w)\xi_i$ where ξ_i is Serre's p -adic zeta pseudomeasure for F_{V_i}/F_{U_i} ([Serre2]). Thus $\xi_i^w = (1-w)\xi_i$.

See [R-W3] Proposition 2 for the more precise proof. $\square$

Let NU_i be the normalizer of U_i in G . Then we have $NU_1 = N\widetilde{U}_2 = NU_3 = G$ and $NU_2 = \widetilde{U}_2$. The quotient group NU_i/U_i acts on the set of locally constant functions on W_i by

$$\varepsilon^\sigma(w) = \varepsilon(\sigma^{-1}w\sigma)$$

where ε is a locally constant functions on W_i and $\sigma \in NU_i/U_i$.

Proposition 7.8 (Sufficient conditions). *Sufficient conditions for the congruences (1), $\widetilde{(2)}$, (2), (3) in Proposition 7.2 to hold are the following (1)', $\widetilde{(2)'}'$, (2)', (3)' respectively:*

- (1)' $\Delta_1^{\varphi(g)}(1-k, \varepsilon) \equiv \Delta_0^g(1-pk, \varepsilon \circ \varphi) \pmod{p\mathbb{Z}_p}$
for all $g \in W_0$ and locally constant $\mathbb{Z}_{(p)}$ -valued function ε on W_1 fixed by $NU_1/U_1 (= G/U_1)$.
- $\widetilde{(2)'}'$ $\widetilde{\Delta}_2^{\varphi(g)}(1-k, \varepsilon) \equiv \Delta_0^g(1-pk, \varepsilon \circ \varphi) \pmod{p\mathbb{Z}_p}$
for all $g \in W_0$ and locally constant $\mathbb{Z}_{(p)}$ -valued function ε on $\widetilde{W}_2$ fixed by $N\widetilde{U}_2/\widetilde{U}_2 (= G/\widetilde{U}_2)$.
- (2)' $\Delta_2^w(1-k, \delta^{(y)}) \equiv 0 \pmod{p\mathbb{Z}_p}$
for all $w \in \Gamma$ and for every coset $y \in W_2/\Gamma^{p^j}$ ($j \in \mathbb{N}$) which is not contained in Γ and fixed by $NU_2/U_2 (= \widetilde{U}_2/U_2)$.
- (3)' $\Delta_3^w(1-k, \delta^{(y)}) \equiv 0 \pmod{p^{m_y}\mathbb{Z}_p}$
for all $w \in \Gamma$ and for every coset $y \in W_3/\Gamma^{p^j}$ ($j \in \mathbb{N}$) which is not contained in Γ . Here p^{m_y} is the order of $(NU_3/U_3)_y = (G/U_3)_y$, the isotropic subgroup of G/U_3 at y .

Proof. The proof of the sufficiency of $\widetilde{(2)'}'$ (resp. (2)') for $\widetilde{(2)}$ (resp. (2)) is completely the same as that of (1)' (resp. (3)') for (1) (resp. (3)). Therefore we only prove the latter two cases.

- The sufficiency of (1)' for (1).

We use similar method to [R-W3], Proposition 4.

The congruence (1) is equivalent to

$$(1 - \varphi(g))\xi_1 \equiv \varphi((1 - g)\xi_0) \pmod{I_1}$$

for an arbitrary $g \in W_0$. Applying the approximation lemma (Proposition 7.7) to them, we obtain

$$(7.3) \quad (1 - \varphi(g))\xi_1 \equiv \sum_{y \in W_1/\Gamma^{p^{j+1}}} \Delta_1^{\varphi(g)}(1 - k, \delta^{(y)})\kappa(y)^{-k}y$$

$$(7.4) \quad \varphi((1 - g)\xi_0) \equiv \sum_{x \in W_0/(W_0^f \times \Gamma^{p^j})} \Delta_0^g(1 - pk, \delta^{(x)})\kappa(x)^{-pk}\varphi(x)$$

$$\text{in } \mathbb{Z}_p[W_1/\Gamma^{p^{j+1}}]/p^{\omega+j+1}$$

$$\text{in } \mathbb{Z}_p[W_0/(W_0^f \times \Gamma^{p^j})]/p^{\omega+j}$$

Note that $\varphi((1 - g)\xi_0)$ is fixed by the conjugate action of G/U_1 since it is contained in $\Lambda(\Gamma)$. $\varphi(g)$ is also fixed by G/U_1 , which implies that $(1 - \varphi(g))\xi_1$ is fixed by G/U_1 (see [R-W3] Lemma 3.2). Therefore we see that $(1 - \varphi(g))\xi_1 - \varphi((1 - g)\xi_0)$ is contained in $(\mathbb{Z}_p[W_1/\Gamma^{p^{j+1}}]/p^{\omega+j+1})^{G/U_1}$ (the G/U_1 -fixed part).

(Case-1) y is fixed by G/U_1 . Then $\delta^{(y)}$ is also fixed by G/U_1 , therefore we may apply (1)' for $\delta^{(y)}$:

$$\Delta_1^{\varphi(g)}(1 - k, \delta^{(y)}) \equiv \Delta_0^g(1 - pk, \delta^{(y)} \circ \varphi) \pmod{p\mathbb{Z}_p}.$$

If $y = \varphi(x) (= \text{Ver}(x))$, x is determined uniquely and $\kappa(y)^{-k} = \kappa(\text{Ver}(x))^{-k} = \kappa(x)^{-pk}$. Moreover, $\delta^{(x)}$ coincides with $\delta^{(y)} \circ \varphi$. This implies that

$$\Delta_1^{\varphi(g)}(1 - k, \delta^{(y)})\kappa(y)^{-k}y - \Delta_0^g(1 - pk, \delta^{(x)})\kappa(x)^{-pk}\varphi(x)$$

$$\in p(\mathbb{Z}_p[W_1/\Gamma^{p^{j+1}}]/p^{\omega+j+1})^{G/U_1} \subseteq I_1^{(j+1)}/p^{\omega+j+1}.$$

If $y \notin \text{Image}(\varphi)$, we have $\delta^{(y)} \circ \varphi = 0$, hence the y -summand vanishes modulo $I_1^{(j+1)}$.

(Case-2) y is not fixed by G/U_1 . Since

$$\Delta_1^{\varphi(g)}(1 - k, \delta^{(y)}) = \Delta_1^{\varphi(g)}(1 - k, \delta^{(y^\sigma)})$$

for every $\sigma \in G/U_1$ (see [R-W3] Lemma 3.2), we have

$$\sum_{\sigma \in G/U_1} (\Delta_1^{\varphi(g)}(1 - k, \delta^{(y)})\kappa(y)^{-k}y)^\sigma$$

$$= \Delta_1^{\varphi(g)}(1 - k, \delta^{(y)})\kappa(y)^{-k} \sum_{\sigma \in G/U_1} y^\sigma \in I_1^{(j+1)}/p^{\omega+j+1}.$$

Therefore (7.3) and (7.4) are congruent modulo $I_1^{(j+1)}/p^{\omega+j+1}$. Taking the projective limit, we obtain the congruence (1) in Proposition 7.2 (Lemma 7.5).

- The sufficiency of (3)' for (3).

Apply the approximation lemma (Proposition 7.7) to $(1 - w)\xi_3$. Then we have

$$(7.5) \quad (1-w)\xi_3 \equiv \sum_{y \in W_3/\Gamma^{p^j}} \Delta_3^w(1-k, \delta^{(y)})\kappa(y)^{-k}y \\ \text{in } \mathbb{Z}_p[W_3/\Gamma^{p^j}]/p^{\omega+j}.$$

Then by (3)',

$$\begin{aligned} & \sum_{\sigma \in (G/U_3)/(G/U_3)_y} \Delta_3^w(1-k, \delta^{(y^\sigma)})\kappa(y^\sigma)^{-k}y^\sigma \\ &= \Delta_3^w(1-k, \delta^{(y)})\kappa(y)^{-k} \sum_{\sigma \in (G/U_3)/(G/U_3)_y} y^\sigma \\ &\equiv p^{m_y} \sum_{\sigma \in (G/U_3)/(G/U_3)_y} \sigma^{-1}y\sigma \quad (\text{by (3)'}) \\ &\in I_3^{(j)}/p^{\omega+j} \end{aligned}$$

for every $y \in W_3/\Gamma^{p^j}$ which is not contained in Γ . Therefore it is obvious that the right hand side of the congruence (7.5) is contained in $(\mathbb{Z}_p[\Gamma/\Gamma^{p^j}] + I_3^{(j)})/p^{\omega+j}$.

By taking the projective limit (Lemma 7.5), we have

$$(1-w)\xi_3 \equiv c_w \pmod{I_3}$$

for a certain element $c_w \in \Lambda(\Gamma)$. Since $1-w \in S_0$, we can obtain the congruence (3) in Proposition 7.2. $\square$

7.4. Hilbert modular forms and Hilbert-Eisenstein series. In this subsection, we review Deligne-Ribet's theory of Hilbert modular forms. See [De-Ri] and [R-W3] Section 3 for more information.

Let K be a totally real number field of degree r , K^∞/K a totally real p -adic Lie extension, and Σ a fixed finite set of primes of K containing all primes which ramify in K^∞ . Let $\mathfrak{f}$ be an integral ideal of $\mathcal{O}_K$ with all prime factors in Σ .

We denote $\mathfrak{h} = \{\tau \in K \otimes \mathbb{C} \mid \text{Im}(\tau) \gg 0\}$ the Hilbert upper-half plane.

For an even positive integer k , we define the action of

$$GL(2, K)^+ = \{g \in GL(2, K) \mid \det(g) \gg 0\}$$

on functions $F: \mathfrak{h} \rightarrow \mathbb{C}$ by

$$(F|_k \begin{pmatrix} a & b \\ c & d \end{pmatrix})(\tau) = \mathcal{N}(ad-bc)^{k/2} \mathcal{N}(c\tau+d)^{-k} F\left(\frac{a\tau+b}{c\tau+d}\right)$$

where $\mathcal{N}: K \otimes \mathbb{C} \rightarrow \mathbb{C}$ is the norm map.

Definition 7.9 (Hilbert modular forms). Let

$$\Gamma_{00}(\mathfrak{f}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, K) \mid a, d \in 1 + \mathfrak{f}, b \in \mathfrak{D}^{-1}, c \in \mathfrak{f}\mathfrak{D} \right\}$$

where $\mathfrak{D}$ is the differential of K . A Hilbert modular form F of weight k on $\Gamma_{00}(\mathfrak{f})$ is a holomorphic function⁹ $F: \mathfrak{h} \rightarrow \mathbb{C}$ satisfying $F|_k M = F$ for all $M \in \Gamma_{00}(\mathfrak{f})$.

We denote the space of Hilbert modular forms of weight k on $\Gamma_{00}(\mathfrak{f})$ by $M_k(\Gamma_{00}(\mathfrak{f}), \mathbb{C})$.

A Hilbert modular form F has a Fourier series expansion (standard q -expansion)

$$c(0) + \sum_{\mu \in \mathcal{O}_K, \mu \gg 0} c(\mu) q_K^\mu$$

where $q_K^\mu = \exp(2\pi\sqrt{-1}\mathrm{Tr}_{K/\mathbb{Q}}(\mu\tau))$.

Deligne and Ribet constructed the Hilbert-Eisenstein series attached to every locally constant function ε on $G = \mathrm{Gal}(K^\infty/K)$ ([De-Ri] Theorem (6.1)).

Theorem-Definition 7.10 (Hilbert-Eisenstein series). *Let ε be a locally constant function on $G = \mathrm{Gal}(K^\infty/K)$ and k a positive even integer. Then there exists an integral ideal $\mathfrak{f} \subseteq \mathcal{O}_K$ with its prime factors in Σ and a Hilbert modular form $G_{k,\varepsilon} \in M_k(\Gamma_{00}(\mathfrak{f}), \mathbb{C})$ whose standard q -expansion is*

$$2^{-r} \zeta_{K^\infty/K}(1-k, \varepsilon) + \sum_{\mu \in \mathcal{O}_K, \mu \gg 0} \left(\sum_{\mu \in \mathfrak{a} \subseteq \mathcal{O}_K, \text{prime to } \Sigma} \varepsilon(\mathfrak{a}) \kappa(\mathfrak{a})^{k-1} \right)$$

Here we denote $\varepsilon(\mathfrak{a}) = \varepsilon\left(\left(\frac{K^\infty/K}{\mathfrak{a}}\right)\right)$ and $\kappa(\mathfrak{a}) = \kappa\left(\left(\frac{K^\infty/K}{\mathfrak{a}}\right)\right)$ where $\left(\frac{K^\infty/K}{\mathfrak{a}}\right)$ is the Artin symbol.

We call $G_{k,\varepsilon}$ the Hilbert-Eisenstein series of weight k attached to ε .

Proof. See [De-Ri] Theorem (6.1). □

Next let us discuss the q -expansion of Hilbert modular forms at cusps. Let $\mathbb{A}_K^f$ be the finite adèle ring of K . By the strong approximation theorem

$$SL(2, \mathbb{A}_K^f) = \hat{\Gamma}_{00}(\mathfrak{f}) \cdot SL(2, K),$$

every $M \in SL(2, \mathbb{A}_K^f)$ is decomposed as $M = M_1 M_2$ where $M_1 \in \hat{\Gamma}_{00}(\mathfrak{f})$ (the closure of $\Gamma_{00}(\mathfrak{f})$ in $SL(2, \mathbb{A}_K^f)$) and $M_2 \in SL(2, K)$. We define $F|_k M$ to be $F|_k M_2$.

For every finite idèle $\alpha \in (\mathbb{A}_K^f)^\times$, set

$$F_\alpha = F|_k \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix}.$$

F_α also has a Fourier series expansion

$$F_\alpha = c(0, \alpha) + \sum_{\mu \in \mathcal{O}_K, \mu \gg 0} c(\mu, \alpha) q_K^\mu,$$

and we call it *the q -expansion of F at the cusp determined by α* .

⁹If $K = \mathbb{Q}$, we also assume that F is holomorphic at ∞ .

Proposition 7.11. *Let k be a positive even integer and ε a locally constant function on G . Then the q -expansion of $G_{k,\varepsilon}$ at the cusp determined by $\alpha \in (\mathbb{A}_K^f)^\times$ is given by*

$$\mathcal{N}((\alpha))^k \left\{ 2^{-r} \zeta_{K^\infty/K}(1-k, \varepsilon_a) + \sum_{\mu \in \mathcal{O}_K, \mu \gg 0} \left(\sum_{\mu \in \mathfrak{a} \subseteq \mathcal{O}_K, \text{prime to } \Sigma} \varepsilon_a(\mathfrak{a}) \kappa(\mathfrak{a})^{k-1} \right) \right\}$$

where (α) is the ideal generated by α and $a = \left(\frac{K^\infty/K}{(\alpha)\alpha^{-1}} \right)$.

Proof. See [De-Ri] Theorem (6.1). $\square$

Now we introduce the Deligne-Ribet's deep principle.

Theorem 7.12 (Deligne-Ribet). *Let F_k ($k \geq 0$) be rational Hilbert modular forms of weight k on $\Gamma_{00}(\mathfrak{f})$ (that is, all coefficients of the q -expansion of F_k at every cusp are rational numbers), and $F_k = 0$ for all but finitely many k .*

Let $\alpha \in (\mathbb{A}_K^f)^\times$. We denote by α_p the p -th component of α . Set

$$S(\alpha) = \sum_{k \geq 0} \mathcal{N} \alpha_p^{-k} F_{k,\alpha}.$$

If $S(\alpha)$ has all coefficients in $p^j \mathbb{Z}_p$ ($j \in \mathbb{Z}$) for one $\alpha \in (\mathbb{A}_K^f)^\times$, $S(\alpha)$ has all coefficients in $p^j \mathbb{Z}_p$ for every $\alpha \in (\mathbb{A}_K^f)^\times$.

Proof. See [De-Ri] Theorem (0.2). $\square$

Corollary 7.13 ([De-Ri] Corollary (0.3)). *Let $S(\alpha)$ as in Theorem 7.12. Suppose that there exists $\alpha \in (\mathbb{A}_K^f)^\times$ such that all non-constant coefficients of $S(\alpha)$ are contained in $p^j \mathbb{Z}_{(p)}$ ($j \in \mathbb{Z}$). Then for arbitrary two distinct elements $\beta, \beta' \in (\mathbb{A}_K^f)^\times$, the difference between the constant terms of the q -expansions of $S(\beta)$ and $S(\beta')$ is an element of $p^j \mathbb{Z}_{(p)}$.*

Proof. Let $c(0, \alpha)$ be the constant term of $S(\alpha)$. We may interpret $c(0, \alpha)$ as an element of $M_0(\Gamma_{00}(\mathfrak{f}), \mathbb{Q})$.

Set $S(\beta)_{c(0,\alpha)} = S(\beta) - c(0, \alpha)$. Then $S(\alpha)_{c(0,\alpha)}$ has all coefficients in $p^j \mathbb{Z}_{(p)}$, so does $S(\beta)_{c(0,\alpha)}$ by Theorem 7.12. Especially the constant term of $S(\beta)_{c(0,\alpha)}$ is also an element of $p^j \mathbb{Z}_{(p)}$, but it is no other than $c(0, \beta) - c(0, \alpha)$. $\square$

7.5. Proof of the sufficient conditions. Now we prove the sufficient conditions in Proposition 7.8. We only prove the conditions (1)' and (3)'. One can prove the conditions $\widetilde{(2)}'$ (resp. (2)') in the same manner as (1)' (resp. (3)').

Condition (1)'. Let k be a positive even integer and ε a locally constant $\mathbb{Z}_{(p)}$ -valued function on W_1 fixed by the action of G/U_1 . Let $G_{k,\varepsilon}$ (resp. $G_{pk,\varepsilon \circ \varphi}$) be the Hilbert-Eisenstein series of weight k (resp. weight pk) attached to ε (resp. $\varepsilon \circ \varphi$) (Theorem-Definition 7.10).

The natural inclusion $F \rightarrow F_{U_1}$ induces the restriction of the Hilbert-Eisenstein series $\text{res } G_{k,\varepsilon}$ on $\mathfrak{h}_F$. It is easy to see that $\text{res } G_{k,\varepsilon}$ is a Hilbert modular form of weight pk and its standard q -expansion is given by ([R-W3] Lemma 7)

$$\begin{aligned} \text{res } G_{k,\varepsilon} &= 2^{-[F_{U_1}:\mathbb{Q}]} \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon) \\ &+ \sum_{\mu \in \mathcal{O}_F, \mu \gg 0} \left(\sum_{(\mathfrak{b}, \nu) \in P_1^\mu} \varepsilon(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} \right) q_F^\mu \end{aligned}$$

where $q_F^\mu = \exp(2\pi\sqrt{-1}\text{Tr}_{F/\mathbb{Q}}(\mu\tau))$ and

$$\begin{aligned} P_1^\mu &= \{(\mathfrak{b}, \nu) \mid \nu \in \mathfrak{b} \subseteq \mathcal{O}_{F_{U_1}}, \mathfrak{b} \text{ is prime to } \Sigma, \\ &\quad \nu \gg 0, \text{Tr}_{F_{U_1}/F}(\nu) = \mu\}. \end{aligned}$$

For $\lambda \in \mathcal{O}_F$, we may construct the Hecke operator U_λ associated to λ (See [R-W3] Lemma 6), which “shifts the coefficients of q -expansion by λ .” Therefore we have

$$\begin{aligned} (\text{res } G_{k,\varepsilon})|_{pk} U_p &= 2^{-[F_{U_1}:\mathbb{Q}]} \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon) \\ &+ \sum_{\mu \in \mathcal{O}_F, \mu \gg 0} \left(\sum_{(\mathfrak{b}, \nu) \in P_1^{p\mu}} \varepsilon(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} \right) q_F^\mu \end{aligned}$$

On the other hand, the standard q -expansion of $G_{pk, \varepsilon \circ \varphi}$ is

$$\begin{aligned} G_{pk, \varepsilon \circ \varphi} &= 2^{-[F:\mathbb{Q}]} \zeta_{F_{V_0}/F_{U_0}}(1-pk, \varepsilon \circ \varphi) \\ &+ \sum_{\mu \in \mathcal{O}_F, \mu \gg 0} \left(\sum_{\mu \in \mathfrak{a} \subseteq \mathcal{O}_F, \text{prime to } \Sigma} \varepsilon \circ \varphi \left(\left(\frac{F_{V_0}/F_{U_0}}{\mathfrak{a}} \right) \right) \kappa(\mathfrak{a})^{pk-1} \right) q_F^\mu. \end{aligned}$$

and note that

$$\varphi \left(\left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) \right) = \text{Ver} \left(\left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{a}} \right) \right) = \left(\frac{F_{V_0}/F_{U_0}}{\mathfrak{a}\mathcal{O}_{F_{U_1}}} \right)$$

by the commutativity of the Artin symbol and the Verlagerung homomorphism (see Lemma 7.4).

Set $S = (\text{res } G_{k,\varepsilon})|_{pk} U_p - G_{pk, \varepsilon \circ \varphi}$, then the μ -th coefficient of S is

$$\sum_{(\mathfrak{b}, \nu) \in P_1^{p\mu}} \varepsilon(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} - \sum_{\mu \in \mathfrak{a} \subseteq \mathcal{O}_F} \varepsilon(\mathfrak{a}\mathcal{O}_{F_{U_1}}) \kappa(\mathfrak{a})^{pk-1}$$

Let $(G/U_1)_{(\mathfrak{b}, \nu)}$ be the isotropic subgroup of G/U_1 at $(\mathfrak{b}, \nu) \in P_1^{p\mu}$.

(Case-1) $(G/U_1)_{(\mathfrak{b}, \nu)} = \{\text{id}\}$. Note that

$$\begin{aligned} \varepsilon \left(\left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{b}^\sigma} \right) \right) &= \varepsilon \left(\sigma^{-1} \left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{b}} \right) \sigma \right) \\ &= \varepsilon^\sigma \left(\left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{b}} \right) \right) \\ &= \varepsilon \left(\left(\frac{F_{V_1}/F_{U_1}}{\mathfrak{b}} \right) \right) \end{aligned}$$

for every $\sigma \in G/U_1$. Similarly we have $\kappa(\mathfrak{b}^\sigma)^{k-1} = \kappa(\mathfrak{b})^{k-1}$. Therefore the sum of $(\mathfrak{b}, \nu)$ -orbit is given by

$$\sum_{\sigma \in G/U_1} \varepsilon(\mathfrak{b}^\sigma) \kappa(\mathfrak{b}^\sigma)^{k-1} = p\varepsilon(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} \in p\mathbb{Z}_{(p)}.$$

(Case-2) $(G/U_1)_{(\mathfrak{b}, \nu)} = G/U_1$. In this case G/U_1 fixes $(\mathfrak{b}, \nu)$, therefore $\nu \in F$ and $\mathfrak{b} = \mathfrak{a}\mathcal{O}_{F_{U_1}}$ for unique integral ideal $\mathfrak{a}$ of F prime to Σ . Since $\text{Tr}_{F_{U_1}/F}(\nu) = p\mu$, we have $\nu = \mu$. Therefore

$$\begin{aligned} \varepsilon(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} &= \varepsilon(\mathfrak{a}\mathcal{O}_{F_{U_1}}) \kappa(\mathfrak{a}\mathcal{O}_{F_{U_1}})^{k-1} = \varepsilon(\mathfrak{a}\mathcal{O}_{F_{U_1}}) \kappa(\mathfrak{a})^{p(k-1)} \\ &\equiv \varepsilon(\mathfrak{a}\mathcal{O}_{F_{U_1}}) \kappa(\mathfrak{a})^{p(k-1)} \pmod{p} \end{aligned}$$

and so the $(\mathfrak{a}\mathcal{O}_F, \mu)$ -term vanishes modulo p .

Therefore S has all the non-constant coefficients in $p\mathbb{Z}_{(p)}$, and we can apply Deligne-Ribet's principle (Corollary 7.13) to $S = S(1)$. There exists a finite idèle γ such that $\left(\frac{F_{V_0}/F_{U_0}}{((\gamma))\gamma^{-1}}\right) = g$ (see [De-Ri] (2.23)). Then by Corollary 7.13 $S(1) - S(\gamma)$ has its constant term in $p\mathbb{Z}_{(p)}$. By easy calculation, the constant term of $E - E(\gamma)$ is

$$\begin{aligned} &2^{-pr} \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon) - 2^{-r} \zeta_{F_{V_0}/F_{U_0}}(1-pk, \varepsilon \circ \varphi) - \mathcal{N}(\gamma_p)^{-pk} \mathcal{N}((\gamma))^{pk} \\ &\quad \times \{2^{-pr} \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon_{\varphi(g)}) - 2^{-r} \zeta_{F_{V_0}/F_{U_0}}(1-pk, (\varepsilon \circ \varphi)_g)\} \\ &= 2^{-pr} \{ \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon) - \kappa(\varphi(g))^k \zeta_{F_{V_1}/F_{U_1}}(1-k, \varepsilon_{\varphi(g)}) \} \\ &\quad - 2^{-r} \{ \zeta_{F_{V_0}/F_{U_0}}(1-pk, \varepsilon \circ \varphi) - \kappa(g)^{pk} \zeta_{F_{V_0}/F_{U_0}}(1-pk, (\varepsilon \circ \varphi)_g) \} \\ &= 2^{-pr} \Delta_1^{\varphi(g)}(1-k, \varepsilon) - 2^{-r} \Delta_0^g(1-pk, \varepsilon \circ \varphi) \\ &\equiv 2^{-r} \{ \Delta_1^{\varphi(g)}(1-k, \varepsilon) - \Delta_0^g(1-pk, \varepsilon \circ \varphi) \} \pmod{p}. \end{aligned}$$

Here we set $r = [F : \mathbb{Q}]$ and use $\mathcal{N}((\gamma)) = \mathcal{N}(\gamma_p) \kappa(g)$ ([R-W3] Lemma 9), $\kappa(\varphi(g)) = \kappa(g)^p$. For precise calculation, see [R-W3], the proof of the Theorem.

Condition (3)'. Let j be a sufficiently large integer and $y \in W_3/\Gamma^{p^j}$ is a coset which is not contained in Γ .

Let $S = G_{k, \delta(y)}$ be the Hilbert-Eisenstein series of weight k attached to $\delta^{(y)}$, then the standard q -expansion of S is given by

$$\begin{aligned} S &= 2^{-[F_{U_3}:\mathbb{Q}]} \zeta_{F_{V_3}/F_{U_3}}(1-k, \delta^{(y)}) \\ &\quad + \sum_{\nu \in \mathcal{O}_{F_{U_3}}, \mu \gg 0} \left(\sum_{\nu \in \mathfrak{b} \subseteq \mathcal{O}_{F_{U_3}}, \text{prime to } \Sigma} \delta^{(y)}(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1} \right) q_{F_{U_3}}^\nu \end{aligned}$$

where $q_{F_{U_3}}^\nu = \exp(2\pi\sqrt{-1}\text{Tr}_{F_{U_3}/\mathbb{Q}}(\nu\tau))$.

(Case-1) $(G/U_3)_{(\mathfrak{b}, \nu)} = \{\text{id}\}$. In this case, we can easily calculate the sum of $(\mathfrak{b}, \nu)$ -orbit :

$$\sum_{\sigma \in G/U_3} \delta^{(y)}(\mathfrak{b}^\sigma) \kappa(\mathfrak{b}^\sigma)^{k-1} = p^{m_y} \sum_{\sigma \in (G/U_3)/(G/U_3)_y} \delta^{(y^{\sigma^{-1}})}(\mathfrak{b}) \kappa(\mathfrak{b})^{k-1}.$$

(Case-2) $(G/U_3)_{(\mathfrak{b},\nu)} \neq \{\text{id}\}$. Let $F_{(\mathfrak{b},\nu)}$ be the fixed field of $(G/U_3)_{(\mathfrak{b},\nu)}$. Then by the same argument as the proof of the condition (1)' (Case-2), there exists $\mu \in F_{(\mathfrak{b},\nu)}$ and $\mathfrak{a} \subseteq \mathcal{O}_{F_{(\mathfrak{b},\nu)}}$ uniquely such that $(\mathfrak{b},\nu) = (\mathfrak{a}\mathcal{O}_{F_{U_3}}, \mu)$. For such $(\mathfrak{a}, \mu)$,

$$\begin{aligned} \delta^{(y)}(\mathfrak{b}) &= \delta^{(y)} \left(\left(\frac{F_{U_3}/F_{V_3}}{\mathfrak{a}\mathcal{O}_{F_{U_3}}} \right) \right) \\ &= \delta^{(y)} \circ \text{Ver} \left(\left(\frac{F_{V_3}/F_{(\mathfrak{b},\nu)}}{\mathfrak{a}} \right) \right) = 0 \end{aligned}$$

because $\text{Im}(\text{Ver}) = \text{Im}(\varphi) \subseteq \Gamma$.

Therefore S has all the non-constant coefficients in $p^{m_y}\mathbb{Z}_{(p)}$. Take a finite idèle γ' such that $\left(\frac{F_{V_3}/F_{U_3}}{(\gamma')^{\gamma'-1}}\right) = w$. Then by Corollary 7.13, the constant term of $S - S(\gamma')$ is also in $p^{m_y}\mathbb{Z}_{(p)}$, and it is equal to

$$2^{-p^3 r} \Delta_3^w (1 - k, \delta^{(y)}).$$

Thus we have finished the proof of Proposition 7.8.

8. PROOF OF THE MAIN THEOREM

Unfortunately, we cannot conclude that $(\xi_i)_i$ is contained in Ψ_S by the congruences obtained in the previous section, so we may not apply Burns' technique (Theorem 3.4) directly to $(\xi_i)_i$.

In this section, we modify the proof of Theorem 3.4 and prove our main theorem (Theorem 3.1) using an inductive technique.

8.1. Kato's p -adic zeta function for F_N/F . Let

$$N = \begin{pmatrix} 1 & 0 & 0 & \mathbb{F}_p \\ 0 & 1 & 0 & \mathbb{F}_p \\ 0 & 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 0 & 1 \end{pmatrix} \times \{1\}$$

be a closed normal subgroup of G and set $\overline{G} = G/N$, $\overline{U}_i = U_i/N$ and $\overline{V}_i = V_i \cdot N/N$.

Then we have the splitting exact sequence

$$1 \longrightarrow N \longrightarrow G \xrightarrow{\pi} \overline{G} \longrightarrow 1.$$

Let

$$s: \overline{G} \rightarrow G; \left(\begin{pmatrix} 1 & a & d \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}, t^z \right) \mapsto \left(\begin{pmatrix} 1 & a & d & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, t^z \right)$$

be a section of π .

The p -adic Lie group

$$\overline{G} = \begin{pmatrix} 1 & \mathbb{F}_p & \mathbb{F}_p \\ 0 & 1 & \mathbb{F}_p \\ 0 & 0 & 1 \end{pmatrix} \times \Gamma$$

is a group “of Heisenberg type,” for which Kazuya Kato has already proven the existence of the p -adic zeta function in [Kato1].

Theorem 8.1 (Kato). *The p -adic zeta function $\bar{\xi}$ for F_N/F exists uniquely and it satisfies the Iwasawa main conjecture (Conjecture 2.15 (2)).*

Sketch of the proof. This theorem is the special case of [Kato1], Theorem 4.1.

First, he constructed the theta map (and its localized version)

$$\begin{aligned}\bar{\theta}: K_1(\Lambda(\bar{G})) &\rightarrow \bar{\Psi} \subseteq \Lambda(\bar{U}_0/\bar{V}_0)^\times \times \Lambda(\bar{U}_1/\bar{V}_1)^\times \\ \bar{\theta}_S: K_1(\Lambda(\bar{G})_S) &\rightarrow \bar{\Psi}_S \subseteq \Lambda(\bar{U}_0/\bar{V}_0)_S^\times \times \Lambda(\bar{U}_1/\bar{V}_1)_S^\times\end{aligned}$$

where $\bar{\Psi}$ (resp. $\bar{\Psi}_S$) was defined to be the subgroup consisting of all elements $(\bar{\eta}_0, \bar{\eta}_1)$ which satisfied the norm relation

$$\mathrm{Nr}_{\Lambda(\bar{U}_0/\bar{V}_0)/\Lambda(\bar{U}_1/\bar{V}_1)} \bar{\eta}_0 \equiv \bar{\eta}_1 \quad (\text{resp. } \mathrm{Nr}_{\Lambda(\bar{U}_0/\bar{V}_0)_S/\Lambda(\bar{U}_1/\bar{V}_1)_S} \bar{\eta}_0 \equiv \bar{\eta}_1)$$

and the congruence

$$\bar{\eta}_1 \equiv \varphi(\bar{\eta}_0) \pmod{\bar{I}_1} \quad (\text{resp. } \pmod{\bar{I}_{S,1}}).$$

In this case, we could show the congruence

$$\bar{\xi}_1 \equiv \varphi(\bar{\xi}_0) \pmod{\bar{I}_{S,1}}$$

directly in the same manner as in §7. Hence, by using Burns’ technique (Theorem 3.4), we might show the existence (and uniqueness) of the p -adic zeta function $\bar{\xi}$ for F_N/F . $\square$

Let $C = C_{F^\infty/F}$ be the complex defined in Definition 2.8. Since we always assume the condition $(*)$ in Proposition 2.10, C is contained in $K_0(\Lambda(G), \Lambda(G)_S)$. Let $[\bar{C}]$ be its image in $K_0(\Lambda(\bar{G}), \Lambda(\bar{G})_S)$. Then Kato’s p -adic zeta function $\bar{\xi}$ satisfies the main conjecture $\partial(\bar{\xi}) = -[\bar{C}]$.

Proposition 8.2. *There exists a characteristic element $f \in K_1(\Lambda(G)_S)$ for F^∞/F whose image in $K_1(\Lambda(\bar{G})_S)$ coincides with $\bar{\xi}$.*

Proof. In the following, we denote by $\bar{x}$ the image of x in $K_1(\Lambda(\bar{G})_S)$ for an element $x \in K_1(\Lambda(G)_S)$.

Let $f' \in K_1(\Lambda(G)_S)$ be a characteristic element of $[C]$. Then by using the functoriality of the connecting homomorphism ∂ , we have $\partial(\bar{\xi} \cdot \bar{f}'^{-1}) = -[\bar{C}] + [\bar{C}] = 0$. By the localization exact sequence (Theorem 1.22), $\bar{u} = \bar{\xi} \cdot \bar{f}'^{-1}$ is the image of an element of $K_1(\Lambda(\bar{G}))$, which we also denote $\bar{u}$ by abuse of notation. Then the element $f = f's(\bar{u})$ satisfies the assertion of the proposition where s denotes the homomorphism $K_1(\Lambda(\bar{G})) \rightarrow K_1(\Lambda(G))$ induced by $s: \bar{G} \rightarrow G$. $\square$

8.2. Completion of the proof. Let $f \in K_1(\Lambda(G)_S)$ be a characteristic element for F^∞/F satisfying Proposition 8.2, that is, $\pi(f) = \bar{\xi}$.

Let $f_i = \theta_{S,i}(f) \in \Lambda(U_i/V_i)_S^\times$ and let $u_i = \xi_i f_i^{-1}$. Then $\partial(u_i) = 0$, so we have $u_i \in \Lambda(U_i/V_i)^\times$ by the localization sequence (Theorem 1.22).

Then it is sufficient to show that $(u_i)_i$ is contained in Ψ : if $(u_i)_i$ is contained in Ψ , there exists $u \in K_1(\Lambda(G))$ such that $\theta_i(u) = u_i$ by the surjectivity of the theta map (Proposition 5.4). One can easily check that $\xi = uf$

satisfies the interpolating properties of the p -adic zeta function for F^∞/F (Definition 2.14) and also satisfies $\partial(\xi) = -[C]$. Namely, ξ is the desired p -adic zeta function.

Note that it was easy to show that $(u_i)_i \in \Psi$ in the proof of Theorem 3.4 because of the assumption $(\xi_i)_i \in \Psi_S$.

Proposition 8.3. *u_i 's satisfy the norm relations in Definition 5.3, and satisfy following congruences:*

$$\begin{aligned} u_1 &\equiv \varphi(u_0) && \text{mod } I_1, \\ \widetilde{u}_2 &\equiv \varphi(u_0) && \text{mod } \widetilde{I}_2, \\ u_2 &\equiv d_2 && \text{mod } I_2'', \\ u_3 &\equiv d_3 && \text{mod } I_3, \end{aligned}$$

where d_2 and d_3 are certain elements of $\Lambda(\Gamma)$.

Proof. f_i 's and ξ_i 's satisfy the norm relations, so do u_i 's.

Since $(f_i)_i$ is contained in Ψ_S , f_i 's satisfy the congruences in Proposition 6.2. On the other hand, ξ_i 's satisfy the congruences in Proposition 7.2. Hence, we can easily show that u_i 's satisfy the desired congruences.¹⁰ $\square$

Lemma 8.4. $(u_i)_i \in \text{Ker} \left(\pi^\times : \prod_i \Lambda(U_i/V_i)^\times \longrightarrow \prod_i \Lambda(\overline{U_i}/\overline{V_i})^\times \right)$.

Proof. This lemma follows from the construction of f and the commutativity of π and norm maps of K -groups (Proposition 1.9). $\square$

Lemma 8.5. $s \circ \pi(I_i) \subseteq I_i$.

Proof. Just simple calculation. $\square$

Lemma 8.6. For each i , $\varphi(u_i) = 1$.

Proof. The Frobenius homomorphism $\varphi: \Lambda(U_i/V_i) \rightarrow \Lambda(\Gamma)$ factors as

$$\begin{array}{ccc} \Lambda(U_i/V_i) & \xrightarrow{\varphi} & \Lambda(\Gamma) \\ \pi \downarrow & & \uparrow s \\ \Lambda(\overline{U_i}/\overline{V_i}) & \xrightarrow{\overline{\varphi}} & \Lambda(\Gamma) \end{array}$$

where $\overline{\varphi}: \Lambda(\overline{U_i}/\overline{V_i}) \rightarrow \Lambda(\Gamma)$ is the Frobenius homomorphism (mod N).

Then this lemma holds since $(u_i)_i \in \text{Ker}(\pi^\times)$ by Lemma 8.4. $\square$

Proof of Theorem 3.1. By operating $s \circ \pi$ to the congruences of Proposition 8.3, we have

$$\begin{aligned} s \circ \pi(u_2) &\equiv d_2 && \text{mod } s \circ \pi(I_2''), \\ s \circ \pi(u_3) &\equiv d_3 && \text{mod } s \circ \pi(I_3). \end{aligned}$$

By Lemma 8.4, both of $s \circ \pi(u_2)$ and $s \circ \pi(u_3)$ are 1. Therefore we obtain

$$(8.1) \quad u_2 \equiv d_2 \equiv 1 = \varphi(u_2) \quad \text{mod } I_2'',$$

$$(8.2) \quad u_3 \equiv d_3 \equiv 1 = \varphi(u_3) \quad \text{mod } I_3,$$

¹⁰Note that we may replace c_2 and c_3 in Proposition 7.2 by invertible elements.

by Lemma 8.5 and 8.6. Hence if we show that

$$(8.3) \quad u_2 \equiv 1 (= \varphi(u_2)) \pmod{I_2},$$

we have $(u_i)_i \in \Psi$, which is the desired result.

Now we show the congruence (8.3). We know that $u_2 \in 1 + I_2''$ and $u_3 \in 1 + I_3$ by (8.1) and (8.2). Note that the p -adic logarithmic homomorphism induces an isomorphism of abelian groups $1 + I_2'' \xrightarrow{\cong} I_2''$ by the same argument as the proof of Lemma 5.6. By the logarithmic isomorphisms (Lemma 5.6), we have $\log u_2 \in I_2''$ and $\log u_3 \in I_3$.

We may describe $\log u_2$ and $\log u_3$ as $\Lambda(\Gamma)$ -linear combinations of generators of I_2'' and I_3 :

$$\begin{aligned} \log u_2 &= \sum_{b,c \neq 0} \tilde{\nu}_{bc}^{(3)} \beta^b \gamma^c h_\zeta + \sum_{b,f} p \tilde{\nu}_{bf}^{(4)} \beta^b \zeta^f, \\ \log u_3 &= \sum_f p^3 \sigma_f^{(1)} \zeta^f + \sum_{e \neq 0} p^2 \sigma_e^{(2)} \varepsilon^e h_\zeta + \sum_{c \neq 0} p \sigma_c^{(3)} \gamma^c h_\varepsilon h_\zeta. \end{aligned}$$

Then we have

$$\mathrm{Tr}_{\Lambda(U_2/V_2)/\Lambda(U_3/V_2)} \log u_2 = \sum_{c \neq 0} p \tilde{\nu}_{0c}^{(3)} \gamma^c h_\zeta + \sum_f p^2 \tilde{\nu}_{0f}^{(4)} h_\zeta$$

and

$$\log u_3 \equiv \sum_f p^2 \left(p \sigma_f^{(1)} + \sum_{e \neq 0} \sigma_e^{(2)} \right) \zeta^f + \sum_{c \neq 0} p^2 \sigma_c^{(3)} \gamma^c h_\zeta \pmod{V_2}.$$

By comparing the coefficients, we have

$$\begin{aligned} \tilde{\nu}_{0f}^{(4)} &= p \sigma_f^{(1)} + \sum_{e \neq 0} \sigma_e^{(2)}, \\ \tilde{\nu}_{0c}^{(3)} &= p \sigma_c^{(3)} \quad (c \neq 0). \end{aligned}$$

Therefore if we set

$$\begin{aligned} \nu_f^{(1)} &= \sigma_f^{(1)}, & \nu_c^{(2)} &= \begin{cases} \sum_{e \neq 0} \sigma_e^{(2)} & \text{if } c = 0, \\ \sigma_c^{(3)} & \text{if } c \neq 0, \end{cases} \\ \nu_{bc}^{(3)} &= \tilde{\nu}_{bc}^{(3)} \quad (b \neq 0, c \neq 0), & \nu_{bf}^{(4)} &= \tilde{\nu}_{bf}^{(4)} \quad (b \neq 0), \end{aligned}$$

we have

$$\begin{aligned} \log u_2 &= \sum_f p^2 \nu_f^{(1)} \zeta^f + \sum_c p \nu_c^{(2)} \gamma^c h_\zeta \\ &\quad + \sum_{b \neq 0, c \neq 0} \nu_{bc}^{(3)} \beta^b \gamma^c h_\zeta + \sum_{b \neq 0, f} p \nu_{bf}^{(4)} \beta^b \zeta^f \end{aligned}$$

which implies $\log u_2 \in I_2$. Hence by the logarithmic isomorphism (Proposition 5.6), u_2 is contained in $1 + I_2$, which implies the congruence (8.3). $\square$

Remark 8.7. By the construction of $\xi = uf$, we have $\theta_S(\xi) = (\xi_i)_i$. Since Ψ_S contains the image of θ_S , we especially obtain the following non-trivial congruences among abelian p -adic pseudomeasures:

$$\begin{aligned}\xi_2 &\equiv \varphi(\xi_0)^p && \text{mod } I_2, \\ \xi_3 &\equiv \varphi(\xi_0)^{p^2} && \text{mod } I_3.\end{aligned}$$

It seems to be impossible to show these congruences directly by using only the theory of Deligne-Ribet.

REFERENCES

- [Bass1] Bass, H., *Algebraic K-theory*, Benjamin (1968).
- [Bass2] Bass, H., *Introduction to some methods of algebraic K-theory*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 20. American Mathematical Society, Providence, R. I. (1974).
- [Ber-Keat] Berrick, A. J., and Keating, M. E., *The localization sequence in K-theory*, *K-Theory* **9** (1995), 577-589.
- [Coates] Coates, J., *p-adic L-functions and Iwasawa's theory*, in *Algebraic Number Fields: L-functions and Galois properties*, Academic Press, London (1977).
- [CFKSV] Coates, J., Fukaya, T., Kato, K., Sujatha, R., and Venjakob, O., *The GL_2 main conjecture for elliptic curves without complex multiplication*, *Publ. Math. Inst. Hautes Études Sci.* **101** (2005), 163-208.
- [De-Ri] Deligne, P., and Ribet, K. A., *Values of abelian L-functions at negative integers over totally real fields*, *Invent. Math.* **59** (1980), 227-286.
- [Fu-Ka] Fukaya, T., and Kato, K., *A formulation of conjectures on p-adic zeta functions in noncommutative Iwasawa theory*, *Amer. Math. Soc. Transl. Ser. 2*, **219** (2006)
- [FW] Ferrero, B., and Washington, L. C., *The Iwasawa invariant μ_p vanishes for abelian number fields*, *Ann. of Math.* **109** (1979), 377-395.
- [Grayson] Grayson, D. R., *K-theory and localization of noncommutative rings*, *J. Pure Appl. Algebra* **18** (1980), 125-127. 18F25 (57R65)
- [Ha-Sh] Hachimori, Y., and Sharifi, R. T., *On the failure of pseudo-nullity of Iwasawa modules*, *J. Algebraic Geom.* **14** (2005), 567-591.
- [Kakde] Kakde, M., *Proof of the main conjecture of noncommutative Iwasawa theory for totally real number fields in certain cases*, [arXiv:0802.2272v2](https://arxiv.org/abs/0802.2272v2) [[math.NT](https://arxiv.org/abs/0802.2272v2)].
- [Kato1] Kato, K., *Iwasawa theory of totally real fields for Galois extensions of Heisenberg type*, preprint.
- [Kato2] Kato, K., *K_1 of some non-commutative completed group rings*, *K-Theory*, **34** (2005), 99-140.
- [Milnor] Milnor, J., *Introduction to Algebraic K-theory*, Princeton Univ. Press, Princeton, N.J., (1971).
- [Oliver] Oliver, R., *Whitehead groups of finite groups*, *London Mathematical Society Lecture Note Series* **132** (1988), Cambridge Univ. Press.
- [Oli-Tay] Oliver, R., and Taylor, L. R., *Logarithmic descriptions of Whitehead groups and class groups for p-groups*, *Mem. Amer. Math. Soc.* **76** (1988), no. 392.
- [R-W1] Ritter, J., and Weiss, A., *Toward equivariant Iwasawa theory*, Part I, *Manuscripta Math.* **109** (2002), 131-146, Part II, *Indag. Math. (N.S.)* **15** (2004), 549-572, Part III, *Math. Ann.* **336** (2006), 27-49, Part IV, *Homology, Homotopy Appl.* **7** (2005), 155-171(electronic).
- [R-W2] Ritter, J., and Weiss, A., *Non-abelian pseudomeasures and congruences between abelian Iwasawa L-functions*, *Pure Appl. Math. Q.* **4** (2008), no. 4, part 1, 1085-1106.
- [R-W3] Ritter, J., and Weiss, A., *Congruences between abelian pseudomeasures*, *Math. Res. Lett.* **15** (2008), no. 4, 715-725.
- [R-W4] Ritter, J., and Weiss, A., *Equivariant Iwasawa theory: an example*, *Doc. Math.* **13** (2008), 117-129.

- [Sch-Ven] Schneider, P., and Venjakob, O., *On the codimension of modules over skew power series rings with applications to Iwasawa algebras*, J. Pure Appl. Algebra **204** (2006), 349-367.
- [Serre1] Serre, J.-P., *Représentations linéaires des groupes finis*, Hermann (1967).
- [Serre2] Serre, J.-P., *Sur le résidu de la fonction zêta p -adique d'un corps de nombres*, C. R. Acad. Sci., Paris **287** (1978), série A, 183-188.
- [Sten] Stenström, B., *Rings of Quotients*, Springer-Verlag, New York-Heidelberg (1975).
- [Swan] Swan, R. G., *Algebraic K-theory*, Lecture Notes in Mathematics, **76**, Springer-Verlag, Berlin-New York (1968).
- [Wall] Wall, C. T. C., *Norms of units in group rings*, Proc. London Math. Soc. (3), **29** (1974), 593-632.
- [Wiles] Wiles, A., *The Iwasawa conjecture for totally real fields*, Ann. of Math. Second Ser., **131** (1990), no.3, 493-540.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 8-1
KOMABA 3-CHOME, MEGURO-KU, TOKYO, 153-8914, JAPAN
E-mail address: thara@ms.u-tokyo.ac.jp